



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in healthcare workers: a meta-analysis of randomized clinical trials

| | |
|-------------------------------|--|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2022-065305 |
| Article Type: | Original research |
| Date Submitted by the Author: | 02-Jun-2022 |
| Complete List of Authors: | <p>Hong, Hwanhee; Duke University, Department of Biostatistics and Bioinformatics</p> <p>Friedland, Anne ; Duke Clinical Research Institute</p> <p>Hu, Mengyi; Duke University, Department of Biostatistics and Bioinformatics</p> <p>Anstrom, Kevin J.; University of North Carolina System, Collaborative Studies Coordinating Center</p> <p>Halabi, Susan; Duke University School of Medicine, Department of Biostatistics & Bioinformatics</p> <p>McKinnon, John; Henry Ford Hospital, Division of Infectious Diseases</p> <p>Amaravadi, Ravi; University of Pennsylvania</p> <p>Rojas-Serrano, Jorge; Instituto Nacional de Enfermedades Respiratorias, Interstitial Lung Disease and Rheumatology Units</p> <p>Abella , Benjamin ; University of Pennsylvania</p> <p>Portillo-Vázquez, Angélica Margarita; Instituto Nacional de Enfermedades Respiratorias</p> <p>Woods, Christopher; Duke Clinical Research Institute</p> <p>Hernandez, Adrian F.; Duke University School of Medicine</p> <p>Boulware, David R.; University of Minnesota Twin Cities</p> <p>Naggie, Susanna; Duke Clinical Research Institute, Infectious Disease</p> <p>Rajasingham, Radha; University of Minnesota Twin Cities</p> |
| Keywords: | COVID-19, STATISTICS & RESEARCH METHODS, EPIDEMIOLOGY |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in healthcare workers: a meta-analysis of randomized clinical trials

Hwanhee Hong, PhD (0000-0002-3736-6327), Anne Friedland, MD (0000-0002-9735-4838), Mengyi Hu, MB (0000-0003-1999-6672), Kevin J. Anstrom, PhD (0000-0001-6452-2172), Susan Halabi, PhD (0000-0003-4135-2777), John E. McKinnon, M.D., M.Sc (0000-0003-3916-4021), Ravi Amaravadi, MD (0000-0002-5768-2474), Jorge Rojas Serrano, MD, PhD (0000-0001-6980-7898), Benjamin S. Abella, MD MPhil (0000-0003-2521-0891), Angélica Margarita Portillo-Vázquez, MD, MSc (0000-0002-5032-4076), Christopher W. Woods, MD, MPH, Adrian Hernandez, MD, MHS (0000-0003-3387-9616), David R Boulware, MD MPH (0000-0002-4715-0060), Susanna Naggie, MD, MHS (0000-0001-7721-6975), Radha Rajasingham, MD (0000-0001-5531-0231)

Hwanhee Hong, Assistant Professor, Department of Biostatistics and Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA

Anne Friedland, Fellow, Department of Infectious Disease, Duke University School of Medicine, 315 Trent Drive, Hanes House, Room 256, Durham, NC 27710, USA

Mengyi Hu, Statistician, Department of Biostatistics and Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA

Kevin J. Anstrom, Director, Collaborative Studies Coordinating Center, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA

Susan Halabi, Professor, Department of Biostatistics and Bioinformatics, Duke University, 2424 Erwin Road, Suite 11088, Durham, NC, 27705, USA

1
2
3 23 John E. McKinnon, Senior Staff, Infectious Diseases, Henry Ford Health, Clara Ford Pavilion,
4
5 24 Suite 322, 2799 W. Grand Blvd., Detroit, MI 48202, USA
6
7 25 Ravi Amaravadi, Professor of Medicine, University of Pennsylvania , 852 BRB 2/3 421 Curie
8
9 26 Blvd., Philadelphia, PA 19063, USA
10
11
12 27 Jorge Rojas Serrano, Attending physician, Interstitial Lung Disease and Rheumatology Unit,
13
14 28 Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas, Calzada de Tlalpan
15
16 29 4502, Tlalpan, Ciudad de México, México. 14080
17
18
19 30 Benjamin S. Abella, Professor, Department of Emergency Medicine, University of Pennsylvania,
20
21 31 423 Guardian Drive Room 412, Philadelphia, PA 19104, USA
22
23 32 Angélica Margarita Portillo-Vázquez, Attending physician, Otolaryngology department, Instituto
24
25 33 Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas. Calzada de Tlalpan 4502,
26
27 34 Tlalpan, Ciudad de México, México. 14080
28
29
30 35 Christopher W. Woods, Professor of Medicine, Duke University, 310 Trent Drive, Durham, NC
31
32 36 27710, USA
33
34
35 37 Adrian Hernandez, Professor of Medicine, Duke Clinical Research Institute, Duke University
36
37 38 300 W. Morgan Street Durham, NC 27701, USA
38
39
40 39 David R Boulware, Professor, Division of Infectious Diseases & International Medicine,
41
42 40 Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis, MN, 55455,
43
44 41 USA
45
46 42 Susanna Naggie, Professor of Medicine, Duke Clinical Research Institute, 300 West Morgan
47
48 43 Street, 5th Floor, Durham, NC 27710, USA
49
50
51
52
53
54
55
56
57
58
59
60

44 Radha Rajasingham, Assistant Professor, Division of Infectious Diseases & International
45 Medicine, Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis,
46 MN, 55455, USA

47
48 **Corresponding author:** Hwanhee Hong, Assistant Professor, Department of Biostatistics and
49 Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA,
50 hwanhee.hong@duke.edu

51
52 Abstract word count: 222

53 Word count: 2258

54
55 **Key words:** SARS-CoV-2, COVID-19, hydroxychloroquine, prophylaxis, health care workers,
56 meta-analysis, clinical trials

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective: We studied the safety and efficacy of hydroxychloroquine (HCQ) as pre-exposure prophylaxis for COVID-19 in healthcare workers (HCWs), using a meta-analysis of randomized controlled trials.

Data Sources: PubMed, EMBASE, EBSCO, and Cochrane databases were searched to identify randomized trials studying HCQ.

Study Selection: Five randomized controlled trials (RCTs) were identified (n=3,672 participants).

Data Extraction and Synthesis: The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis between HCQ and placebo using a Bayesian random-effects model. A *pre-hoc* statistical analysis plan was written, and the review protocol was registered at PROSPERO (CRD42021285093)

Main Outcomes: The primary efficacy outcome was polymerase chain reaction (PCR)-confirmed SARS-CoV-2 infection and the primary safety outcome was incidence of adverse events. The secondary outcome included clinically suspected SARS-CoV-2 infection.

Results: Compared with placebo, HCWs randomized to hydroxychloroquine (HCQ) had no significant difference in PCR-confirmed SARS-CoV-2 infection (odds ratio [OR] 0.60, 95% credible interval [CI]: 0.24, 1.28), clinically suspected SARS-CoV-2 infection (OR 0.76, 95% CI: 0.48, 1.24), or adverse events (OR 1.46, 95% CI: 0.87, 2.22).

Conclusions and Relevance: Our meta-analysis of five RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in HCWs found that HCQ does not significantly reduce the risk of confirmed or clinically suspected SARS-CoV-2 infection or significantly increase adverse events compared with placebo.

90 INTRODUCTION

91 Early during the SARS-CoV-2 pandemic, based on *in vitro* antiviral activity of both chloroquine
92 and hydroxychloroquine against SARS-CoV-2 ¹⁻³, clinicians considered use of
93 hydroxychloroquine (HCQ) for treatment and prevention of SARS-CoV-2 infection and the
94 associated disease, COVID-19. While there are now published randomized controlled trials of
95 HCQ for the treatment of COVID-19 in the inpatient and outpatient setting ^{4,5}, there remains a
96 lack of adequately powered randomized controlled trials of HCQ for the pre-exposure
97 prophylaxis (PrEP) of SARS-CoV-2 infection. A number of PrEP studies were planned early in
98 the pandemic; however, several never opened to enrollment and those that did open were closed
99 early without reaching full accrual due to the rapidly changing landscape of preventative
100 therapies, including vaccines, and a significant shift in public opinion of HCQ as a medical
101 intervention for SARS-CoV-2.

102 Yet, as vaccination access remains insufficient globally ⁶, studying the pre-exposure
103 prophylaxis potential for a drug with a known safety profile is crucial to protect people at high
104 risk of exposures, such as healthcare workers (HCWs) ^{7,8}. Two large randomized, placebo-
105 controlled trials testing the safety and efficacy of HCQ as pre-exposure prophylaxis for COVID-
106 19 in HCWs, PATCH ⁹ and Minnesota (MN)-COVID-PREP ¹⁰, showed potential for a modest
107 benefit of HCQ but were both underpowered, if a modest effect exists. In addition, more trials ¹¹⁻
108 ¹³ studying HCQ as pre-exposure prophylaxis of COVID-19 in HCWs have since been
109 completed and with similar limitations.

110 To address the most common limitation, inadequate power to show a modest effect, we
111 conducted a formal meta-analysis of pre-exposure prophylactic HCQ studies in HCWs. We
112 conducted a systematic search for clinical trials of pre-exposure prophylactic use of HCQ against

infection of SARS-CoV-2 in HCWs, thoroughly compared similarities and differences in characteristics of the identified studies and performed a Bayesian meta-analysis to combine results of the trials.

METHODS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis¹⁴. A statistical analysis plan was written in advance and the review protocol was registered at PROSPERO (CRD42021285093).

Search strategy and information sources

We searched PubMed/Medline, Ovid/Embase, EBSCO/CINAHL, and Cochrane databases from database inception through the final search date October 11, 2021. We used keywords related to COVID-19, HCQ, and prophylaxis. The full search strategies are provided in eTable 1. Unpublished data from eligible randomized controlled trials listed on ClinicalTrials.gov and other relevant information were obtained by contacting the study authors and principal investigators.

Eligibility criteria and study selection

The eligibility criteria included phase II or phase III randomized controlled trials (RCTs) of hydroxychloroquine for use as pre-exposure prophylaxis in HCWs with moderate to high risk of exposure. We excluded observational studies, crossover trials, studies where the method of allocation to treatment was not truly random, duplicate studies, and non-original data studies. No language, publication date, or publication status restrictions were applied. References of prior

systematic reviews and meta-analyses were also screened for related studies. Study selection involved screening of titles and abstracts followed by full-text evaluation of possible eligible studies.

Data collection process

Each of the selected studies were independently reviewed by two reviewers (AF, MH, or HH). We extracted data on the study design, baseline characteristics, interventions, and outcomes. Any disagreements of collected information between reviews were reconciled through discussion by all three reviewers.

Outcome measures

The primary efficacy outcome for the meta-analysis was laboratory confirmed SARS-CoV-2 infection by polymerase chain reaction (PCR) test and the primary safety outcome was incidence of adverse events (Table 1). The secondary efficacy outcome was suspected or probable SARS-CoV-2 infection. Included studies had the following outcome definitions: (1) laboratory confirmed SARS-CoV-2 infection defined as COVID-19 like symptoms and positive SARS-CoV-2 PCR and (2) suspected or probable SARS-CoV-2 infection defined as COVID-19 like symptoms but lack of confirmatory PCR testing.

Table 1. Treatment strategies, adherence, trial-defined primary outcome, and follow-up time in each trial

| Trial (NCT ID) | Trial-defined primary outcome | Follow-up | Treatment group | Randomized treatment assignment | Randomized sample size |
|-------------------------------|--|-----------|-----------------|--|------------------------|
| HERO-HCQ (NCT04334148) | Confirmed (by NP swab PCR) or suspected COVID-19 infection | 60 days | HCQ | HCQ 600 mg BID loading dose for Day 1, followed by 400 mg QD for 29 days | 683 |

| | | | | | |
|---|-------------------|-----------|------------------|----------------------|-----|
| | through 30 days | | Placebo | Placebo | 676 |
| PATCH (NCT04329923) | COVID-19 | 56 days | HCQ | HCQ 600mg daily for | 64 |
| | infection as | (8 weeks) | | 60 days | |
| | determined by | | Placebo | Placebo | 61 |
| | positive NP swab | | | | |
| | over 8 weeks | | | | |
| MN-COVID- PREP (NCT04328467) | COVID-19 free | 84 days | HCQ ¹ | HCQ loading doses | 989 |
| | survival time by | (12 | | (400mg twice 6-8hrs | |
| | lab confirmed or | weeks) | | apart) followed by | |
| | probable illness | | | 400mg once weekly | |
| | | | | or 400mg twice | |
| | | | | weekly for 84 days | |
| | | | Placebo | Placebo | 494 |
| Rojas-Serrano et al. (NCT04318015) | Time to | 60 days | HCQ | HCQ 200 mg daily | 62 |
| | symptomatic | | | for 60 days | |
| | respiratory | | Placebo | Placebo | 65 |
| | infection with a | | | | |
| | positive COVID | | | | |
| | RT PCR over 60 | | | | |
| | days | | | | |
| WHIP (NCT04341441) | Lab confirmed | 56 days | HCQ ² | HCQ 400 mg loading | 387 |
| | cases of COVID- | (8 weeks) | | dose for Day 1, | |
| | 19 determined by | | | followed by 200 mg | |
| | either IgM and | | | daily or 400 mg | |
| | IgG serology in | | | weekly on the same | |
| | blood sample or | | | day of each week for | |
| | RT-PCR test | | | 56 days | |
| | results Confirmed | | Placebo | Placebo | 191 |
| | new cases of | | | | |
| | COVID-19 | | | | |

HCQ=Hydroxychloroquine

¹ HCQ group in the MN-COVID-PREP study includes participants taking 400 mg once weekly or 400 mg twice weekly.

² HCQ group in the WHIP study includes participants taking 200 mg daily or 400 mg weekly.

Treatment assignment

Our meta-analysis did not study HCQ dosing specific effects. For studies randomizing participants to more than one HCQ arm with different doses, all HCQ arms were merged and considered as a single HCQ arm. For example, the MN-COVID-PREP and WHIP studies each had HCQ arms with weekly or twice weekly dosing, thus these two arms were combined as a single HCQ arm for the meta-analysis (Table 1).

Risk of bias within individual studies

172 Two independent reviewers (AF, HH) assessed the risk of bias (low, intermediate, high) of the
173 included studies using the Cochrane's Collaboration tool ¹⁵ (eTable 2).

175 **Statistical analysis**

176 Bayesian logistic regression meta-analysis models under two assumptions (fixed effect and
177 random effects) were fitted to estimate the odds ratio of having an outcome between
178 hydroxychloroquine and placebo ¹⁶. The fixed effect model assumes that the odds ratio is
179 constant across studies, while the random effects model accounts for heterogeneity in the odds
180 ratios across studies. To assess and compare the goodness-of-fit of the fitted fixed and random
181 effects models, we calculated the Watanabe-Akaike information criterion ¹⁷. In the Bayesian
182 models, we assigned non-informative prior distributions as no prior information was available.
183 The odds ratios and the associated 95% confidence intervals were estimated using Markov chain
184 Monte Carlo (MCMC) algorithms. In addition, we calculated Bayesian posterior probabilities of
185 the odds ratio smaller than 1 or 0.5 for the primary efficacy outcome, and greater than 2 for the
186 safety outcome ¹⁸. The standard deviation of the random effects and I^2 ¹⁹ were estimated to
187 quantify the between-study heterogeneity, where small values of both metrics indicate little
188 heterogeneity. All analyses were conducted using the `rstan` package (version 2.21.2)²⁰ in R
189 4.0.2 ²¹. We used two parallel chains, where each chain consists of 50,000 samples after a
190 25,000-sample burn-in. We checked convergence of the MCMC chains for all model parameters
191 using trace plots and Gelman-Rubin diagnostic statistics ²².

193 **Patient and public involvement**

194 No patient involved.

RESULTS

Search results

Our database search resulted in 164 unique studies after excluding duplicates. Of those, 161 studies were screened out due to irrelevance based on title and abstract screening. Three studies were assessed in full-text for eligibility and they met the inclusion criteria (Figure 1). Of those, two trials, conducted by the University of Pennsylvania (NCT04329923, denoted by PATCH)⁹ and the University of Minnesota (NCT04328467, denoted by MN-COVID-PREP)¹⁰, recruited healthcare workers (HCWs) while the third cluster-randomized trial, conducted by the National University of Singapore (NCT04446104), recruited non-HCWs²³ was excluded from the meta-analysis. Additionally, we identified three eligible trials via ClinicalTrials.gov that were completed but had not yet been published in peer-reviewed journals and included in the meta-analysis. These three studies recruited HCWs and were conducted by Duke University (NCT04334148, denoted by HERO-HCQ)¹¹, the National Institute of Respiratory Diseases of Mexico (NCT04318015, denoted by Rojas-Serrano et al.)¹², and the Henry Ford Health System (NCT04341441, denoted by WHIP)¹³. As a result, a total of five studies in a population consisting of HCWs were identified. The secondary efficacy outcome of suspected or probable SARS-CoV-2 infection was reported by HERO-HCQ, MN-COVID-PREP, and WHIP studies.

Study and patient characteristics

Study design, population, treatment strategies, and key characteristics are presented in Table 1 and eTable 3. A total of 3,672 randomized participants (2,185 randomized to HCQ) from the 5 studies were included in the meta-analysis. The five studies defined HCWs broadly and included first responders (emergency medical services, fire, and police). The follow-up duration of the 5

studies ranged from 56 days to 84 days. The HCQ dosing scheme varied across studies, including daily dosing ranging from 200 to 600mg daily with or without a loading dose and once or twice weekly dosing. The duration of therapy also varied across studies with a range of 30 to 84 days (Table 1). The trial-specific definitions of primary outcome and adverse events are comparable across trials (Table 1, eTable 4).

Baseline characteristics by randomized treatment assignment are reported (eTable 5). The HERO-HCQ, MN-COVID-PREP, and WHIP studies had average age between 40 and 45, while PATCH and Rojas-Serrano et al. included relatively younger participants with average age between 31 and 34 years. The aggregate proportion of women within each study varied across the 5 trials, with a range from 51% to 69%. In addition, the PATCH and Rojas-Serrano et al. studies had smaller sample size compared with the other three studies and showed a difference in female ratio between placebo and HCQ groups. In the HERO-HCQ, PATCH, MN-COVID-PREP, and WHIP studies, over 80% of study participants were white. The PATCH and MN-COVID-PREP studies had high proportions of HCWs working in an emergency department (56% and 41%, respectively) and the PATCH study had a high proportion of nurses (67%).

Treatment adherence was assessed by two methods, self-reported adherence and/or pill count at the end of the study. MN-COVID-PREP additionally conducted remote blood sampling to verify HCQ concentrations in a subset. Adherence varied significantly across the studies, with a low proportion of approximately 52% in the Rojas-Serrano et al. study and 97-98% in the PATCH study.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results of meta-analysis

Overall, 1.2% (45/3672) developed PCR-confirmed SARS-CoV-2 infection and 5.8% (200/3420) developed suspected COVID-19 that was not laboratory confirmed. Since the goodness-of-fit assessment using Watanabe-Akaike information criterion concluded that the random effects meta-analysis model was as good as or better than the fixed effect meta-analysis model for all outcomes, we reported the results under the random effects model. Compared with placebo, HCWs randomized to HCQ had numerically lower rate of PCR-confirmed SARS-CoV-2 infection cases (odds ratio [OR] 0.60, 95% credible interval [CI]: 0.24, 1.28), and suspected or probable SARS-CoV-2 infection cases (OR 0.76, 95% CI: 0.48, 1.24). Participants treated with HCQ had a numerically higher rate of adverse events (OR 1.46, 95% CI: 0.87, 2.22) (Figure 2). None of these odds ratios were statistically significant. The outcome data used in our analyses are presented in eTable 6.

The Bayesian posterior probabilities of the odds ratio less than 1 for the confirmed SARS-CoV-2 infection outcome (i.e., the probability of HCQ favoring over placebo) was 0.92, while the posterior probability of odds ratio less than 0.5 (i.e., the probability that the odds of having a confirmed SARS-CoV-2 infection outcome in HCQ is less than a half of the odds in placebo) was 0.32. The posterior probability of the odds ratio greater than 2 for the adverse event outcome (i.e., the probability that the odds of having an adverse event in HCQ is greater than twice of the odds in placebo) was 0.05.

Our meta-analysis showed little or moderate variability of effect estimates across studies with I^2 value of 0%, 0%, and 55%, and the estimated standard deviation of the random effects of 0.30,

0.25, and 0.38 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively.

DISCUSSION

Understanding the pre-exposure prophylactic effect of HCQ against COVID-19 remains relevant, as its use continues, particularly in the international setting^{24 25}. Our meta-analysis of the five RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in HCWs found that HCQ did not have a statistical association with fewer confirmed or suspected/probable SARS-CoV-2 infection cases compared with placebo. While the odds ratios of the studies and the meta-analysis all favor HCQ, the confidence intervals remain wide suggesting low confidence in the true point estimate. Furthermore, in this population, COVID-19 events rates were low, particularly for the most relevant PCR-confirmed infection outcome. The low event rate raises further concern for the uncertainty of these outcomes. Thus, if there is a minimal effect, the absolute benefit would be low. To gain more certainty, a very large study would need to be done and this is difficult to support now due to availability of highly effective vaccines. The safety profile of HCQ in the outpatient setting is well understood²⁶. In these outpatient studies there was no significant difference in adverse events in the HCQ versus the placebo arm, confirming the well-known safety profile of HCQ.

Our findings can be applied to HCWs but should not be generalized to a broader population. Our systematic search found only one published RCT of pre-exposure prophylaxis from Singapore that was not in HCW. This study showed a significant reduction in the risk of COVID-19 infection in the HCQ arm when compared with the comparator arm, vitamin C. However, this

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

study showed moderate risk of bias as it used an open-label cluster-randomization design, the Institutional Review Board excluded higher risk persons from the hydroxychloroquine arm only, and the participants may not be representative of a general population due to the communal living environment.

A prior meta-analysis ²⁷ investigated pre-exposure (two RCTs included) and post-exposure (three RCTs included) prophylactic effects of HCQ and found insignificant effects on SARS-CoV-2 infection and adverse events, similar to ours. For the pre-exposure prophylactic effects, our meta-analysis includes three additional RCTs, resulting in the most up-to-date, systematic, and comprehensive evidence.

Although a meta-analysis allows for combining evidence from multiple studies in a principled way, our meta-analysis has limitations. First, our analysis did not evaluate effects of different HCQ doses and combined two weekly dosing HCQ arms using different doses in each of MN-COVID-PREP and WHIP studies. The five RCTs included in our meta-analysis studied five different dosing schemes and a meta-analysis using aggregate-level data is not a sufficient source to study dosing effects. Second, detailed subgroup analyses were not conducted due to limited information. Individual-level data are required to study both dosing and subgroup effects.

Our meta-analysis of five RCTs investigating safety and efficacy of HCQ as pre-exposure prophylaxis in HCWs provides the most up-to-date evidence on HCQ. We found that HCQ does not reduce the risk of confirmed or probable SARS-CoV-2 infection or adverse events compared

with placebo. Hydroxychloroquine should not be used for pre-exposure prophylaxis in the HCW population.

Contributors

All authors fulfill the ICMJE criteria for authorship. Drs. Hong, Naggie, Rajasingham, and Anstrom designed the study. Drs. Hong and Friedland and MS Hu collected and analyzed the data. Drs. Hong, Naggie, and Rajasingham wrote the manuscript and all authors provided critical review. All authors approved and decided to submit the paper for publication.

Competing interests

All authors except Dr. Abella reported no financial relationship with commercial interest. Dr. Abella have received NIH funds for COVID-19 related research, and holds equity in VOC Health, a start-up company that is developing novel covid testing.

Funding

This study is funded by the Patient Centered Outcomes Research Institute (PCORI), Contract Number COVID-19-2020-001. The funder had no role in the design, conduct, analysis, or reporting of this study.

Data sharing statement

The data are presented in eTable 6.

REFERENCES

1
2
3 333 1. Kalil AC. Treating COVID-19—off-label drug use, compassionate use, and randomized
4
5 334 clinical trials during pandemics. *JAMA* 2020;323(19):1897-98.
6
7
8 335 2. McCreary EK, Pogue JM, Pharmacists obotSoID. Coronavirus Disease 2019 Treatment: A
9
10 336 Review of Early and Emerging Options. *Open Forum Infectious Diseases* 2020;7(4) doi:
11
12 337 10.1093/ofid/ofaa105
13
14
15 338 3. Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently
16
17 339 emerged novel coronavirus (2019-nCoV) in vitro. *Cell research* 2020;30(3):269-71.
18
19 340 4. RECOVERY Collaborative Group. Effect of hydroxychloroquine in hospitalized patients with
20
21 341 Covid-19. *New England Journal of Medicine* 2020;383(21):2030-40.
22
23
24 342 5. Skipper CP, Pastick KA, Engen NW, et al. Hydroxychloroquine in nonhospitalized adults with
25
26 343 early COVID-19: a randomized trial. *Annals of internal medicine* 2020;173(8):623-31.
27
28 344 6. Padma T. COVID vaccines to reach poorest countries in 2023—despite recent pledges. *Nature*
29
30 345 2021;595(7867):342-43.
31
32
33 346 7. World Health Organization. Prevention, identification and management of health worker
34
35 347 infection in the context of COVID-19 2020 [Available from:
36
37 348 <https://www.who.int/publications/i/item/10665-336265> accessed May 13th 2022.
38
39
40 349 8. The United Kingdom Office for National Statistics. Coronavirus (COVID-19) infections in the
41
42 350 community in England: May 2021 2021 [Available from:
43
44 351 [https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsa](https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristics)
45
46 352 [nddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristic](https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristics)
47
48 353 [sofpeopletestingpositiveforcovid19incountriesoftheuk20may2021#percentage-testing-](https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristics)
49
50 354 [positive-for-covid-19-by-patient-facing-and-non-patient-facing-job-roles-uk](https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristics) accessed
51
52 355 May 13th 2022.
53
54
55
56
57
58
59
60

9. Abella BS, Jolkovsky EL, Biney BT, et al. Efficacy and safety of hydroxychloroquine vs placebo for pre-exposure SARS-CoV-2 prophylaxis among health care workers: a randomized clinical trial. *JAMA internal medicine* 2021;181(2):195-202.
10. Rajasingham R, Bangdiwala AS, Nicol MR, et al. Hydroxychloroquine as Pre-exposure Prophylaxis for Coronavirus Disease 2019 (COVID-19) in Healthcare Workers: A Randomized Trial. *Clinical Infectious Diseases* 2020;72(11):e835-e43. doi: 10.1093/cid/ciaa1571
11. Naggie S, Milstone A, Castro M, et al. Hydroxychloroquine for pre-exposure prophylaxis of COVID-19 in health care workers: a randomized, multicenter, placebo-controlled trial (HERO-HCQ). *medRxiv* 2021
12. Rojas-Serrano J, Thirion AMP-VI, Vázquez-Pérez J, et al. Hydroxychloroquine For Prophylaxis Of COVID-19 In Health Workers: A Randomized Clinical Trial. *medRxiv* 2021
13. McKinnon JE, Wang DD, Zervos M, et al. Safety and tolerability of hydroxychloroquine in health care workers and first responders for the prevention of COVID-19: WHIP COVID-19 Study. *International Journal of Infectious Diseases* 2022;116:167-73.
14. Hutton B, Salanti G, Caldwell DM, et al. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Annals of internal medicine* 2015;162(11):777-84.
15. Sterne JA, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *bmj* 2019;366
16. Hong H, Wang C, Rosner GL. Meta-analysis of rare adverse events in randomized clinical trials: Bayesian and frequentist methods. *Clinical Trials* 2021;18(1):3-16.

1
2
3 379 17. Watanabe S, Oppen M. Asymptotic equivalence of Bayes cross validation and widely
4
5 380 applicable information criterion in singular learning theory. *Journal of machine learning*
6
7 381 *research* 2010;11(12)
8
9
10 382 18. Ferreira D, Ludes P-O, Diemunsch P, et al. Bayesian predictive probabilities: a good way to
11
12 383 monitor clinical trials. *British journal of anaesthesia* 2021;126(2):550-55.
13
14 384 19. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in*
15
16 385 *medicine* 2002;21(11):1539-58.
17
18
19 386 20. Stan Development Team. 2020. RStan: the R interface to Stan. R package version 2.21.2.
20
21 387 <http://mc-stan.org/>
22
23 388 21. R Core Team. 2014. R: A Language and Environment for Statistical Computing.
24
25 389 <https://www.R-project.org/>
26
27 390 22. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences.
28
29 391 *Statistical science* 1992;7(4):457-72.
30
31 392 23. Seet RCS, Quek AML, Ooi DSQ, et al. Positive impact of oral hydroxychloroquine and
32
33 393 povidone-iodine throat spray for COVID-19 prophylaxis: An open-label randomized trial.
34
35 394 *International Journal of Infectious Diseases* 2021;106:314-22.
36
37 395 24. Infante M, Ricordi C, Alejandro R, et al. Hydroxychloroquine in the COVID-19 pandemic
38
39 396 era: in pursuit of a rational use for prophylaxis of SARS-CoV-2 infection. *Expert review*
40
41 397 *of anti-infective therapy* 2021;19(1):5-16.
42
43 398 25. Revised advisory on the use of hydroxychloroquine (HCQ) as prophylaxis for SARS-CoV-2
44
45 399 infection (in supersession of previous advisory dated 23rd March. 2020): Indian Council
46
47 400 of Medical Research; 2022 [Available from:

- 1
2
3 401 [https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ](https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ_SARS_CoV2_infection.pdf)
4
5 402 [SARS CoV2 infection.pdf](https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ_SARS_CoV2_infection.pdf).
6
7
8 403 26. Safety of hydroxychloroquine among outpatient clinical trial participants for COVID-19.
9
10 404 Open forum infectious diseases; 2020. Oxford University Press US.
11
12 405 27. García-Albéniz X, Amo Jd, Polo R, et al. Systematic review and meta-analysis of
13
14 406 randomized trials of hydroxychloroquine for the prevention of COVID-19. *medRxiv*
15
16
17 407 2021:2020.09.29.20203869. doi: 10.1101/2020.09.29.20203869
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. Flowchart of literature review

Figure 2. Forest plots of the meta-analysis results showing the number of events (y), sample size (n), posterior median of odds ratios, and the associated 95% credible intervals comparing HCQ versus placebo for (a) lab-confirmed positive COVID-19, (b) suspected COVID-19, and (c) adverse events.

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

Identification

Screening

Eligibility

Included

| | |
|---------------------------|------------|
| Initial Systematic Search | |
| PubMed | 141 |
| Embase | 39 |
| EBSCO | 0 |
| Cochrane | 6 |
| TOTAL | 186 |

Article Screened
N=164

Full-text articles accessed for eligibility
N=3

Studies including healthcare workers
N=5

Duplicates Excluded **N=22**:

- Across database (N=5)
- Within database (N=17)


Full-text articles excluded, with reasons **N=161**:

- Study on Post-exposure of COVID (N=52)
- Protocol (N=17)
- Reviews (N=39)
- Retracted (N=2)
- Non-randomized (N=9)
- No RCT involved (N=42)

Study on non-healthcare workers excluded
N=1

Unpublished trials on healthcare workers included
N=3


| HCQ y/n | Placebo y/n | Odds Ratio [95% CI] |
|----------------|----------------|--------------------------|
| 3/683 | 6/676 | 0.49 [0.12, 1.98] |
| 4/64 | 4/61 | 0.95 [0.23, 3.98] |
| 11/989 | 6/494 | 0.91 [0.34, 2.49] |
| 1/62 | 6/65 | 0.16 [0.02, 1.38] |
| 2/387 | 2/191 | 0.49 [0.07, 3.51] |
| 21/2185 | 24/1487 | 0.60 [0.24, 1.28] |


$$I^2 = 0\%$$

Favors Hydroxychloroquine Favors Placebo

Odds ratio for confirmed positive COVID-19


| HCQ y/n | Placebo y/n | Odds Ratio [95% CI] |
|----------------|----------------|--------------------------|
| 38/683 | 47/676 | 0.79 [0.51, 1.23] |
| 47/989 | 33/494 | 0.70 [0.44, 1.10] |
| 22/387 | 13/191 | 0.83 [0.41, 1.68] |
| 07/2059 | 93/1361 | 0.76 [0.48, 1.24] |


$$I^2 = 0\%$$

Favors Hydroxychloroquine Favors Placebo

Odds ratio for suspected COVID-19

| HCC y/n | Placebo y/n | Odds Ratio [95% CI] |
|----------------|-----------------|--------------------------|
| 16/683 | 13/676 | 1.22 [0.58, 2.56] |
| 29/65 | 17/65 | 2.27 [1.09, 4.76] |
| 16/989 | 100/494 | 1.85 [1.43, 2.39] |
| 32/62 | 38/65 | 0.76 [0.38, 1.53] |
| 92/387 | 85/191 | 1.23 [0.87, 1.74] |
| 35/2186 | 253/1491 | 1.46 [0.87, 2.22] |


$$I^2 = 55\%$$

For Favors Hydroxychloroquine Favors Placebo
Odds ratio for adverse event

Supplementary Materials

CONTENTS

- eTable 1. Search code
- eTable 2. Risk of bias
- eTable 3. Characteristics of included trials
- eTable 4. Definition of adverse events
- eTable 5. Baseline characteristics
- eTable 6. Results of outcome measures in each study

For peer review only

eTable 1. Search code that was used to identify publications as of October 11, 2021

PubMed search

| | |
|----|--|
| #1 | ((covid[Title/Abstract]) OR (coronavirus[Title/Abstract])) OR (sars-cov[Title/Abstract]) |
| #2 | (hcq[Title/Abstract]) OR (hydroxychloro[Title/Abstract]) |
| #3 | (prophyl[Title/Abstract]) OR (Prep[Title/Abstract]) |
| #4 | (randomized clinical trial[Publication Type]) OR (controlled clinical trial[Publication Type]) OR (randomized[Title/Abstract]) OR (randomised[Title/Abstract]) |
| #5 | #1 AND #2 AND #3 AND #4 |

Embase search

| | |
|----|--|
| #1 | covid:ab,ti OR coronavirus:ab,ti OR 'sars cov':ab,ti |
| #2 | prep:ab,ti OR prophylaxis:ab,t |
| #3 | 'randomized controlled trial':ab,ti OR 'randomized clinical trial':ab,ti |
| #4 | hydroxychloroquine:ab,ti OR hcq:ab,ti |
| #5 | #1 AND #2 AND #3 AND #4 |


























Ebsco search

| | |
|----|--|
| S1 | TX covid OR TX coronavirus OR TX sars-cov |
| S2 | TX hydroxychloroquine OR TX HCQ |
| S3 | TX prep OR TX prophyl |
| S4 | TX randomized clinical trial OR TX controlled clinical trial |
| S5 | S1 AND S2 AND S3 AND S4 |

Cochrane search

| | |
|----|--|
| #1 | (covid):ti,ab,kw OR (coronavirus):ti,ab,kw OR ("SARS-CoV"):ti,ab,kw (Word variations have been searched) |
| #2 | ("hydroxychloroquine"):ti,ab,kw OR (hcq):ti,ab,kw |
| #3 | (prophyl):ti,ab,kw OR (prep):ti,ab,kw |
| #4 | ("randomized clinical trial"):pt OR (controlled clinical trial):pt OR (randomized):ti,ab,kw OR (randomised):ti,ab,kw |
| #5 | #1 AND #2 AND #3 |
| #6 | #4 AND #5 |

eTable 2. Risk of bias of included trials using the Cochrane risk assessment tool. Green circle is for low risk and yellow circle is for some concerns

| | HERO-HCQ NCT04334148 | PATCH NCT04329923 | MN-COVID- PREP NCT04328467 | Rojas-Serrano et al. NCT04318015 | WHIP NCT04341441 |
|---|---|---|---|---|---|
| Selection bias (Randomization process) |  |  |  |  |  |
| Performance bias (Deviations from the intended interventions) |  |  |  |  |  |
| Attrition bias ¹ (Missing outcome data) |  |  |  |  |  |
| Reporting bias (Measurement of the outcome) |  |  |  |  |  |
| Other sources of bias (Selection of the reported result) |  |  |  |  |  |

¹ All studies but the Mexico study reported minimal loss to follow-up (<10%). The Mexico study reported 18% (25/130) lost to follow-up and additional 12% (16/130) discontinued the intervention.

eTable 3. Characteristics of included trials

| | HERO-HCQ NCT04334148 | PATCH NCT04329923 | MN-COVID-PREP NCT04328467 | Rojas-Serrano et al. NCT04318015 | WHIP NCT04341441 |
|---|--|--|--|---|--|
| N (randomization) | 1360 | 132 | 1496 | 130 | 624 |
| Study start date ¹ | 4/22/2020 | 4/9/2020 | 4/6/2020 | 4/21/2020 | 4/10/2020 |
| Study completion date ² | 1/9/2021 | 11/13/2020 | 7/13/2020 | 3/31/2021 | 12/14/2020 |
| Occupation | HCWs at risk of COVID exposure through work in the ICU, emergency department, emergency services, respiratory services or COVID unit | HCWs (Physicians, nurses, certified nursing assistants, emergency technicians, respiratory therapists) eligible working >20 hrs/week | HCWs (physicians, nurses, emergency medical technicians) with direct contact with COVID patients including emergency department and ICU setting, first responders and performing aerosol generating procedures | HCWs (nurses, nursing aids, cleaning staff, orderlies, respiratory therapists and physicians) taking care of hospitalized patients with COVID | HCW, first responders and correlational/law officers, nursing home workers, medical students, public transit workers, household family members of HCW in Michigan and Ohio |
| Sites | 34 sites across the US | 2 tertiary urban hospitals | Multiple sites nationwide across US and Canada | Single site (National Institute of Respiratory Diseases of Mexico) | Multiple sites at Michigan in the US |
| Randomization | Yes (Phase III) | Yes (Phase II) | Yes (Phase III) | Yes (Phase III) | Yes (Phase III) |
| Trial type | Double-blinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | No weight requirement | <40kg excluded | <50kg excluded | N/A |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Excluded | Excluded | Excluded | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Included | Exclude |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Included | |
| Significant renal disease (stage IV or greater) | Excluded | Included | Excluded | Excluded | Exclude |
| Pregnant/breastfeeding | Included | Excluded | Included in US, Excluded in Canada | Excluded | Exclude |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Excluded | Included | Exclude |
| Use of other medications with significant drug interactions | Included | Excluded | Excluded | Included | N/A |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine or azithromycin) | Any treatment for COVID-19 within 14 days excluded | Current use of HCQ or chloroquine excluded | HCQ or chloroquine within 30 days excluded | Chronic use of HCQ included |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | N/A | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Excluded | Excluded if symptoms within 2 weeks unless negative COVID test | Excluded | Excluded | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | N/A |
| Positive COVID serology | Included | Included | N/A | Included | N/A |
| Analysis | Modified intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat |

HCW=Healthcare workers; ICU=Intensive care unit; ¹ Date when first participant was enrolled; ² Date when final data were collected for the last participant

eTable 4. Definition of adverse events

| RCT | AE definition |
|---|---|
| HERO-HCQ NCT04334148 | Adverse events include general disorders and administration site conditions, psychiatric disorders, skin and subcutaneous tissue disorders, cardiac disorders, infections and infestations, nervous system disorders, gastrointestinal disorders, investigations (electrocardiogram QT prolonged and heart rate increased), ear and labyrinth disorders, renal and urinary disorders, and respiratory, thoracic and mediastinal disorders. |
| PATCH NCT04329923 | Adverse events include abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. |
| MN-COVID-PREP NCT04328467 | Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and others. |
| Rojas-Serrano et al. NCT04318015 | Examples of adverse events are as follows: abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and other. |
| WHIP NCT04341441 | Covid-19 related symptoms, covid-19 clinical disease and medication adverse effects including gastrointestinal disorders, nervous system disorders, respiratory, thoracic and mediastinal disorders, general disorders and administration site conditions, cardiac disorders, musculoskeletal and connective tissue disorders, psychiatric disorders, skin and subcutaneous tissue disorders, ear and labyrinth disorders, and eye disorders. |

eTable 5. Baseline characteristics with additional variables and detailed information. Sample mean and standard deviation (in parenthesis) are reported for continuous variables, and the number of participants and proportion (in parenthesis) are reported for binary or categorical variables.

| | | HERO-HCQ NCT04334148 | | PATCH NCT04329923 | | MN-COVID-PREP NCT04328467 | | Rojas-Serrano et al. NCT04318015 | | WHIP NCT04341441 | |
|---|---------------------------------------|-------------------------|--------------------|-------------------------|-------------------------|------------------------------|--------------------------|-------------------------------------|-------------------------------|--|-------------|
| | | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo | HCQ | Placebo | HCQ ² | Placebo |
| N (ITT) | | 683 | 676 | 66 | 66 | 989 | 494 | 62 | 65 | 387 | 191 |
| Age | | 44.2 (11.9) | 43.1 (11.2) | 31 (20-66) ³ | 34 (23-62) ³ | 41.5 (35, 49) ³ | 40 (34, 48) ³ | 31.0 (26.4-39) ⁴ | 31.9 (27.2-43.7) ⁴ | 45.7 (11.6); 44.9 (11.4) ² | 44.1 (12.7) |
| Female | | 442 (64.7%) | 446 (66.0%) | 54 (82%) | 37 (56%) | 519 (52.5%) | 241 (48.8%) | 29 (42.6%) | 42 (64.6%) | 220 (57%) | 114 (60%) |
| BMI (kg/m^2) | | 28.3 (6.3) | 28.6 (6.7) | 26 (19-37) ⁵ | 26 (20-50) ⁵ | | | 26.7 (3.9) | 27.2 (4.6) | | |
| Current smoker | | | | 0 (0%) | 0 (0%) | 38 (3.84%) | 13 (2.6%) | 20 (32.2%) ⁶ | 23 (35.4%) ⁶ | | |
| Race/ Ethnicity | White | 624 (91.4%) | 610 (90.2%) | 55 (83%) | 54 (82%) | 852 (86.1%) | 419 (84.8%) | | | 334 (86%) | 161 (84%) |
| | Asian | | | 7 (11%) | 7 (11%) | 46 (4.7%) | 29 (5.9%) | | | 23 (6%) | 15 (8%) |
| | African American | 18 (2.6%) | 23 (3.4%) | 3 (4%) | 1 (2%) | 10 (1.0%) | 10 (2.0%) | | | 15 (4%) | 9 (5%) |
| | Hispanic | 39 (5.7%) | 40 (5.9%) | 0 (0%) | 2 (3%) | 40 (4.0%) | 18 (3.6%) | | | 11 (3%) | 7 (4%) |
| Comorb idities | Asthma | 58 (8.5%) | 77 (11.4%) | 9 (14%) | 14 (21%) | 91 (9.2%) | 59 (11.9%) | | | | |
| | Diabetes | 20 (2.9%) | 35 (5.2%) | 1 (2%) | 3 (5%) | 36 (3.6%) | 14 (2.8%) | | | | |
| | Hypertension | 99 (14.5%) | 99 (14.6%) | 3 (5%) | 14 (21%) | 145 (14.7%) | 60 (12.1%) | | | | |
| | None | | | 54 (82%) | 40 (61%) | 646 (65.3%) | 336 (68.0%) | 53 (85.5%) | 58 (89.2%) | | |
| Practice Location | Emergency Department | 96 (14.1%) | 94 (13.9%) | 38 (58%) | 36 (55%) | 417 (42.2%) | 190 (38.5%) | | | 48 (12%) | 19 (10%) |
| | Internal Medicine ward | | | 17 (26%) | 18 (27%) | 98 (9.9%) | 56 (11.3%) | | | 31 (8%) | 20 (10%) |
| | ICU/anesthesia | | | 6 (9%) | 6 (9%) | | | | | | |
| | Labor and delivery | | | 5 (7%) | 6 (9%) | | | | | | |
| | Ambulance | 66 (9.7%) | 63 (9.3%) | | | 73 (7.4%) | 45 (9.1%) | | | | |
| | Congregate care setting | | | | | 46 (4.7%) | 20 (4.0%) | | | | |
| | ICU | 48 (7.0%) | 59 (8.7%) | | | 184 (18.6%) | 85 (17.2%) | | | 37 (10%) | 23 (12%) |
| | Operating room | | | | | 103 (10.4%) | 75 (15.2%) | | | | |
| Occupation | EMS, Fire and Police First Responders | | | | | | | | | 32 (8%) | 16 (8%) |
| | Nurse | 186/677 (27.5%) | 167/668 (25.0%) | 46 (70%) | 42 (64%) | | | | | | |
| | Physician | 143/677 (21.1%) | 144/668 (21.6%) | 11 (17%) | 16 (24%) | | | | | | |
| | Certified Nurse Assistant | | | 2 (3%) | 2 (3%) | | | | | | |
| | ED Technician | | | 3 (4%) | 1 (2%) | | | | | | |
| | Respiratory therapist | 15/677 (2.2%) | 18/668 (2.7%) | 3 (4%) | 5 (7%) | | | | | | |
| | Nurse or Physician | | | | | | | 31 (50%) | 33 (50.8%) | | |
| | Emergency Medicine Provider | | | | | 407 (41.1%) | 190 (38.5%) | | | | |
| | ICU provider | | | | | 160 (16.2%) | 83 (16.8%) | | | | |
| | Anesthesia/ENT | | | | | 178 (18.0%) | 105 (21.3%) | | | | |
| HCW in COVID unit | | | | | | 76 (7.7%) | 29 (5.9%) | | | | |
| Healthcare worker in congregated care setting | | | | | | 11 (1.1%) | 4 (0.8%) | | | | |
| First responder | | | | | | 115 (11.6%) | 65 (13.2%) | | | | |

1 HCQ=Hydroxychloroquine ; ITT= Intention-to-treat ; BMI=Body mass index ; ICU=Intensive care unit; ED=Emergency department ; ENT=Ear, nose, throat ; HCW=Healthcare
2 worker

3 ¹ HCQ group in the MN-COVID-PREP study included participants taking 400 mg once weekly or 400 mg twice weekly.

4 ² HCQ group in the WHIP study included participants taking 200 mg daily or 400 mg weekly.

5 ³ Median (range)

6 ⁴ Median (IQR)

7 ⁵ Mean (range)

8 ⁶ Current or previous smoker

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

For peer review only

eTable 6. Results of outcome measures in each study. Sample size and the number of participants who had each outcome are reported with proportions (%) in parentheses.

| | HERO-HCQ NCT04334148 | | PATCH NCT04329923 | | MN COVID-PREP NCT04328467 | | Rojas-Serrano et al. NCT04318015 | | WHIP NCT04341441 | |
|--|-------------------------|----------|----------------------|-----------|------------------------------|------------|-------------------------------------|-----------|---------------------|-----------|
| Treatment | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo | HCQ | Placebo | HCQ ² | Placebo |
| N (ITT) | 683 | 676 | 64 | 61 | 989 | 494 | 62 | 65 | 387 | 191 |
| Primary Outcome | | | | | | | | | | |
| Confirmed COVID-19 | 3 (0.4) | 6 (0.9) | 4 (6.3) | 4 (6.6) | 11 (1.1) | 6 (1.2) | 1 (1.6) | 6 (9.2) | 2 (0.5) | 2 (1.0) |
| Suspected with COVID compatible symptoms | 38 (5.6) | 47 (7.0) | | | 47 (4.8) | 33 (6.7) | | | 22 (5.7) | 13 (6.8) |
| Secondary outcome | | | | | | | | | | |
| Adverse event ³ | 16 (2.3) | 13 (1.9) | 29 (45.3) | 17 (27.9) | 316 (32.0) | 100 (20.2) | 32 (51.6) | 38 (58.5) | 192 (49.6) | 85 (44.5) |

HCQ= Hydroxychloroquine; ITT= Intention-to-treat; AE=Adverse event ; COVID-RS=COVID-19 related symptoms ; Vit C= Vitamin C
¹ HCQ group in the MN-COVID-PREP study included participants taking 400 mg once weekly or 400 mg twice weekly.
² HCQ group in the WHIP study included participants taking 200 mg daily or 400 mg weekly.
³ Number of patients with any adverse events



PRISMA 2020 Checklist

| Section and Topic ! | Item # | Checklist item ! | Location where item is reported ! |
|---------------------------------------|--------|--|-----------------------------------|
| TITLE ! | | | ! |
| "\$&!! | '! | ()&\$*+,!\$-&!&./O.\$!12!1!2,2\$&3 1\$#4!&.5\$&67! | '! |
| ABSTRACT ! | | | ! |
| 892\$.14\$!! | :! | ; &&!\$- &!<=(; >8!: ? : ?!+O.!892\$.14\$2!4- &4@#2\$7! | A! |
| INTRODUCTION ! | | | ! |
| =1\$#0*1%!! | A! | B&24.#9&\$- &!&.1\$#0*1%!!+O.!\$- &!&.5\$&6!#*\$!\$- &!40*\$&C\$!O+!&C#2\$*#D!@*06%&)D&7! | E! |
| F9&4\$5&2!! | H! | <.05#&) &!1*!&C/!#4\$!2\$1\$&3 &*\$!O+!\$- &!O9&4\$5&5&12!O!.!KL&2\$0*!2J!\$- &!&.5\$&6!1)).&22&27! | M! |
| METHODS ! | | | ! |
| N#D#9#\$, !4.\$&.#1!! | E! | ; /&4#+,!\$- &!#*4%L2#0*!1*)!&C4%L2#0*!4.\$&.#1!+O.!\$- &!&.5\$&6!1*)!-06!2\$!)#&2!6&.&!D.O!/&)!+O.!\$- &!2,*\$- &2&27! | O! |
| (*0.3 1\$#0*! 20L.4&2!! | M! | ; /&4#+,!1%!)1\$1912&2P!&.D#2\$&.2P!6&92\$&2P!O.D1*#21\$#0*2P!&.&.&.*4&!#2\$2!1*)!O\$- &!20L.4&2!2&1.4- &!)O.!40*2L\$&!)\$O!#&) &*\$#+,!2\$!)#&27!; /&4#+,!\$- &!&1\$&!6- &*\$!&14-!20L.4&!6!2!!12\$!2&1.4- &!)O.!40*2L\$&!)7! | M! |
| &1.4-!2\$.1\$&D,!&1&4\$#0*!/.04&22! | O! | <.&2&*\$!\$- &!+L%2&1.4-!2\$.1\$&D#&2!+O.!1%!)1\$1912&2P!&.D#2\$&.2!1*)!6&92\$&2P!#*4%L)#*D!1* ,!+L%2!1*)!#3\$2!L2&!)7! | M! |
| &1&4\$#0*!/.04&22! | Q! | ; /&4#+,!\$- &!3&\$-0)2!L2&!)\$O!&#&) &!6- &\$- &.&!1!2\$!) ,!3&\$!\$- &!#*4%L2#0*!4.\$&.#1!O+!\$- &!&.5\$&6P!#*4%L)#*D!-06!3 1* ,!&.5\$&6&.2!24.&.&*&!)&14-!&.40.!)1*)!&14-!&/O.\$!&.&5&!)P!6- &\$- &.&!6O.@&!)#*)&/ &*) &*\$\$,P!1*)!#1! / /%419%&P!)&\$1#2!O+!1L\$0 3 1\$#0*!\$00!2!L2&!)#*!\$- &! / .04&227! | O! |
| B1\$140%&4\$0*! /.04&22!! | R! | ; /&4#+,!\$- &!3&\$-0)2!L2&!)\$O!40%&4\$!)1\$1!+.03!&/O.\$2P!#*4%L)#*D!-06!3 1* ,!&.5\$&6&.2!40%&4\$&!)1\$1!+.03!&14-!&/O.\$P!6- &\$- &.&!6O.@&!)#*)&/ &*) &*\$\$,P!1* ,!/.04&22&2!+O.!O9\$1*#*#D!O.!40*#&.3*#D!)1\$1!+.03!2\$!) ,!#*5&2\$D1\$0.2P!1*)!#1! / /%419%&P!)&\$1#2!O+!1L\$0 3 1\$#0*!\$00!2!L2&!)#*!\$- &! / .04&227! | O! |
| B1\$1!\$&3 2!! | ' ?1! | S#2\$!1*)!&#* &!1%!!O!\$40 3&2!+O.!6-#4-!)1\$1!6&.&!20LD-\$7!; /&4#+,!6- &\$- &.&!1%!!&.2L\$2!\$-1\$!6&.&!40 3 /1\$9&!6\$-!&14-!O!\$40 3&2!)0 3 1*#*!&14-!2\$!) ,!6&.&!20LD-\$!I&7D7!+O.!1%!!3&12L.&2P!\$3&! /O#*\$2P!1*1% ,2&2J!1*)!#*!O\$P!\$- &!3&\$-0)2!L2&!)\$O!&#&) &!6-#4-!&.2L\$2!\$O!40%&4\$7! | OTQ! |
| | ' ?9! | S#2\$!1*)!&#* &!1%!!O!\$- &!5!&.19!&2!+O.!6-#4-!)1\$1!6&.&!20LD-\$!I&7D7!/1.\$#4/1*1!1*)!#* &\$.5&*\$#0*!4-1.14\$&.#2\$42P!L*)#*D!20L.4&27!B&24.#9&!1* ,!122L 3 /\$#0*2!3 1)!&190L\$1* ,!3&22*#D!O.!L*4&!1.#*+0.3 1\$#0*7! | OTQ! |
| \$L) ,!#2@!O+!9#12! 122&22 3&*\$! | ' !! | ; /&4#+,!\$- &!3&\$-0)2!L2&!)\$O!122&22!&.2@!O+!9#12!#*\$- &!#*4%L)&!)2\$!)#&2P!#*4%L)#*D!)&\$1#2!O+!\$- &!\$00!2!J!L2&!)P!-06!3 1* ,!&.5\$&6&.2!122&22&!)&14-!2\$!) ,!1*)!6- &\$- &.&!\$- &.&!6O.@&!)#*)&/ &*) &*\$\$,P!1*)!#1! / /%419%&P!)&\$1#2!O+!1L\$0 3 1\$#0*!\$00!2!L2&!)#*!\$- &! / .04&227! | Q! |
| N+&4\$! 3&12L.&2!! | ' :! | ; /&4#+,!+O.!&14-!O!\$40 3&\$- &!&+&4\$! 3&12L.&2J!I&7D7!&.2@!1.\$#O!P! 3&1*)!#* &.*4&J!L2&!)#*!\$- &!2,*\$- &2#2!O! / .&2&*\$1\$#0*!O+!&.2L\$27! | QTR! |
| 2 ; ,*\$- &2#2! 3&\$-0)2! | ' A! | B&24.#9&\$- &! / .04&22&2!L2&!)\$O!&#&) &!6-#4-!2\$!)#&2!6&.&!&#D#9&!+O.!&14-!2,*\$- &2#2!I&7D7!\$19L!1\$*#D!\$- &!2\$!) ,!#* &\$.5&*\$#0*!4-1.14\$&.#2\$42!1*)! | ; L / /%3 &*\$! |
| | ' A9! | B&24.#9&!1* ,!3&\$-0)2!&.KL#&!)\$O! / .&/1.&!\$- &!&)1\$1!+O! / .&2&*\$1\$#0*!O!2,*\$- &2#2!2L4-!12!-1*)#*D!O+!3&22*#D!2L 3 3 1 ,!2\$1\$2\$42P!O!)!1\$! | QTR! |
| | ' A4! | B&24.#9&!1* ,!3&\$-0)2!L2&!)\$O!\$19L!1\$&!O!5#2L1%!)#2/!1 ,!&.2L\$2!O!#*)#5#&L1!2\$!)#&2!1*)!2,*\$- &2&27! | QTR! |
| | ' A)! | B&24.#9&!1* ,!3&\$-0)2!L2&!)\$O!2,*\$- &2#V&!&.2L\$2!1*)! / .05#&) &!1!1.\$#0*1%!!+O.!\$- &!4-0#4!2J7!(+!3&\$!T1*1% ,2#2!6!2! /&.&+0.3&#&P!)&24.#9&\$- &!30)&!2J!P! 3&\$-0)2!J!\$O!#&) &*\$#+,!\$- &! / .&2&4&!1*)!&C\$&*\$!O+!2\$1\$2\$41!- &\$&.0D&*&#\$,P!1*)!20+6 1.&! /14@1D&2J!L2&!)7! | QTR! |
| | ' A&! | B&24.#9&!1* ,!3&\$-0)2!L2&!)\$O!&C/!O.&! /022#9&!41L2&2!O+!- &\$&.0D&*&#\$,!1 3 0*#D!2\$!) ,!&.2L\$2!I&7D7!2L9D.O! /!1*1% ,2#2P! 3&\$!T.&D.&22#0*J7! | QTR! |
| | ' A+! | B&24.#9&!1* ,!2&*2\$5\$,!1*1% ,2&2!40*)L4\$&!)\$O!122&22!&.O9L2\$*#22!O+!\$- &!2,*\$- &2#V&!)&.2L\$27! | QTR! |
| =&/O.\$*#D!9#12! 122&22 3&*\$! | ' H! | B&24.#9&!1* ,!3&\$-0)2!L2&!)\$O!122&22!&.2@!O+!9#12!)L&!\$O!3&22*#D!&.2L\$2!#*1!2,*\$- &2#2!1.2#*D!+.03!&/O.\$*#D!9#12&2J7! | ; L / /%3 &*\$! |
| W&.\$1#*\$,!&122&22 3&*\$! | ' E! | B&24.#9&!1* ,!3&\$-0)2!L2&!)\$O!122&22!&.4&.\$1#*\$,!O!&.40*#&!)&4&J!#*\$- &!90) ,!O+!&5#&) &4&+O.!1*!O!\$40 3&7! | ; L / /%3 &*\$! |



PRISMA 2020 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

| Section and Topic ! | Item # | Checklist item ! | Location where item is reported ! |
|--|--------|---|-----------------------------------|
| RESULTS ! | | | ! |
| ; \$L) , !2&1&4\$0*!! | 'M1! | B&24.#9&1\$- &!&2L\$2!0!\$- &!2&1.4-1!*)!2&1&4\$0*!/ .04&22!+.03!\$- &!*L39&.!0!&.40.)2!#)&*\$#&!)#*!\$- &!2&1.4-!\$0!\$- &!*L39&.!0!2\$L)&2!#*4!L)&)*!!\$- &!&.5&6!#)&1% , !L2#*D!1!+!06!)*1D.137! | RT' ?! |
| | 'M9! | W\$&12\$L)&2!\$-1\$!3D-\$!1//&1.!\$0!3&&\$!\$- &!#*4!L2#0*!4.#\$&.1P!9L\$!6-#4-!6&.&!&C4!L)&)*!1*)!&C/!1#*!6- , !\$- & , !6&.&!&C4!L)&)*7! | RT' ?! |
| ; \$L) , !4-1.14\$&. #2\$42!! | 'O! | W\$&1&14-!#*4!L)&)*2\$L)& , !1*)!/.&2&*\$!\$2!4-1.14\$&. #2\$427! | RT' ' ! |
| =#2@!0!9#12!#*!2\$L)&2!! | 'Q! | <.&2&*\$!122&223&*2!0!&.#2@!0!9#12!+0.!&14-!#*4!L)&)*2\$L)& , 7! | Q! |
| =&2L\$2!0!#*)#5#)L1!2\$L)&2!! | 'R! | X0.!1\$!0L\$403&2P!/.&2&*\$!+0.!&14-!2\$L)& , Y!1!J!2L331. , !2\$1\$2\$42!+0.!&14-!D.0L/!16- &.&1//.0/.#1\$&J!1*)!19J!1*!&+&4\$!&2\$31\$&1!*)!\$2!/.&4#2#0*! !&7D!40*#)&*4&Z4.&)*9!&1#*\$&.51!J!#)&1% , !L2#*D!2\$.L4\$L.&)*!19&2!0!./!\$0\$27! | ; L / /%3&*\$! |
| =&2L\$2!0!2, *\$- &2&2! | : ?1! | X0.!&14-!2, *\$- &2#2P!9.#&+ , !2L331.#2&1\$- &!4-1.14\$&. #2\$42!1*)!&.#2@!0!9#12!130*!D!40*\$.#9L\$#*D!2\$L)&27! | ; L / /%3&*\$! |
| | : ?9! | <.&2&*\$!&2L\$2!0!1\$!2\$1\$2\$41!\$2, *\$- &2&2!40*)L4\$&)*7!(!3&\$1T1*1%, 2#2!6!2!)0* &P!/.&2&*\$!+0.!&14-!\$- &!2L331. , !&2\$31\$&1!*)!\$2!/.&4#2#0*! !&7D!40*#)&*4&Z4.&)*9!&1#*\$&.51!J!1*)!3&12L.&2!0!2\$1\$2\$41!- &.\$&.0D&* &\$, 7!(!403/1.#*D!D.0L/2P!)&24.#9&1\$- &!&.#4\$0*!0!\$- &!&+&4\$7! | ' 'T' : ! |
| | : ?4! | <.&2&*\$!&2L\$2!0!1\$!#*5&2\$D!\$0*2!0!+/022#9!&14!L2&2!0!- &.\$&.0D&* &\$, !130*!D!2\$L)& , !&2L\$27! | ' 'T' : ! |
| | : ?)! | <.&2&*\$!&2L\$2!0!1\$!2&*2\$5\$, !1*1% , 2&2!40*)L4\$&)*!\$0!122&22!\$- &!&.09L2\$* &22!0!\$- &!2, *\$- &2#V&)*!&2L\$27! | ' 'T' : ! |
| =&/0.#\$*D!9#12&2! | : '! | <.&2&*\$!122&223&*2!0!&.#2@!0!9#12!)L&!\$0!3#22#*D!&2L\$2!1!&2#*D!+0.03!&/0.#\$*D!9#12&2J!+0.!&14-!2, *\$- &2#2!122&22&)*7! | ; L / /%3&*\$! |
| W&.#1#*\$, !0!&5#)&*4&!! | : :! | <.&2&*\$!122&223&*2!0!&.#1#*\$, !10.!40*#)&*4&J!#*\$!\$- &!90) , !0!&5#)&*4&!+0.!&14-!0L\$403&!122&22&)*7! | ; L / /%3&*\$! |
| DISCUSSION ! | | | ! |
| B#24L22#0*!! | : A1! | <.05#)&!1!D&* &.1!#*\$&. / .&\$1\$0*!0!\$- &!&2L\$2!#*\$!\$- &!40* &\$&C!0!0\$- &.&!&5#)&*4&7! | ' :T' H! |
| | : A9! | B#24L22!1* , !#3\$1\$0*2!0!\$- &!&5#)&*4&1#*4!L)&)*#*\$!\$- &!&.5&67! | 'AT' H! |
| | : A4! | B#24L22!1* , !#3\$1\$0*2!0!\$- &!&.5&6!/.04&22&2!L2&)*7! | 'AT' H! |
| | : A)! | B#24L22!3 / #41\$0*2!0!\$- &!&2L\$2!+0!/.14\$4&P!/0#4 , P!1*)!+L\$L.&!&2&1.4-7! | 'H! |
| OTHER INFORMATION ! | | | ! |
| =&D#2\$.1\$0*!1*)!/.0\$040!! | : H1! | <.05#)&!&D#2\$.1\$0*!#*+0.31\$0*!+0!\$- &!&.5&6!P!#*4!L)&)*D!&D#2\$&.!*13&1!*)!&D#2\$.1\$0*!+0!L39&.P!0.!2\$1\$&1\$-1\$!\$- &!&.5&6!6!2!+0\$!&D#2\$&.&)*7! | ; L / /%3&*\$! |
| | : H9! | (*)#41\$&16- &.&1\$- &!&.5&6!/.0\$040!41*!9&144&22&)*P!0.!2\$1\$&1\$-1\$!1!/.0\$040!6!2!+0\$!/.&/1.&)*7! | O! |
| | : H4! | B&24.#9&1!*)!&C/!1#*!1* , !13&*)3&*2!\$0!#*+0.31\$0*!/.05#)&)*!1\$!&D#2\$.1\$0*!0!&#*\$!\$- &!/.0\$040!7! | O! |
| ; L / /0.\$! | : E! | B&24.#9&1!20L.&2!0!+!#*1*4#1!0!&1*0*!T#*1*4#1!2L//0.\$!+0!\$- &!&.5&6!P!1*)!\$- &!&0!&10!\$- &!+L*)&2!0!&2/0*20.2!#*\$!\$- &!&.5&67! | A! |
| W03 / &\$#*D!#*\$&. &2\$2! | : M! | B&4!1.&!1* , !403 / &\$#*D!#*\$&. &2\$2!0!&.5&6!1L\$-0.27! | A! |
| 85!#19#&\$, !0!)1\$1P!40)&!1*)!0\$- &!31\$&. #1\$2! | : O! | =&/0.\$!6-#4-!0!\$- &!&0!06#*D!1.&!/L9#4% , !151#19!&1!*)!6- &.&1\$- & , !41*!9&!0L*)Y!&3 / !1\$&!1\$!140%&4\$0*!+0.32!)1\$!1!&C\$.14\$&)*.03!#*4!L)&)*!2\$L)&2!)1\$!1!L2&)*!0!&1!1!1*1% , 2&2!1*1% , \$4!40)&[!1* , !0\$- &!31\$&. #1\$2!L2&)*!#*\$!\$- &!&.5&67! | ; L / /%3&*\$! |

BMJ Open

Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in healthcare workers: a meta-analysis of randomized clinical trials

| | |
|---------------------------------|--|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2022-065305.R1 |
| Article Type: | Original research |
| Date Submitted by the Author: | 03-Apr-2023 |
| Complete List of Authors: | Hong, Hwanhee; Duke University, Department of Biostatistics and Bioinformatics Friedland, Anne; UNC School of Medicine Hu, Mengyi; Duke University, Department of Biostatistics and Bioinformatics Anstrom, Kevin J.; University of North Carolina System, Collaborative Studies Coordinating Center Halabi, Susan; Duke University School of Medicine, Department of Biostatistics & Bioinformatics McKinnon, John; Henry Ford Hospital, Division of Infectious Diseases Amaravadi, Ravi; University of Pennsylvania Rojas-Serrano, Jorge; Instituto Nacional de Enfermedades Respiratorias, Interstitial Lung Disease and Rheumatology Units Abella, Benjamin; University of Pennsylvania Portillo-Vázquez, Angélica Margarita; Instituto Nacional de Enfermedades Respiratorias Woods, Christopher; Duke Clinical Research Institute Hernandez, Adrian F.; Duke University School of Medicine Boulware, David R.; University of Minnesota Twin Cities Naggie, Susanna; Duke Clinical Research Institute, Infectious Disease Rajasingham, Radha; University of Minnesota Twin Cities |
| Primary Subject Heading: | Infectious diseases |
| Secondary Subject Heading: | Infectious diseases, Evidence based practice |
| Keywords: | COVID-19, STATISTICS & RESEARCH METHODS, EPIDEMIOLOGY |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in
healthcare workers: a meta-analysis of randomized clinical trials**

Hwanhee Hong, PhD (0000-0002-3736-6327), Anne Friedland, MD (0000-0002-9735-4838),
Mengyi Hu, MB (0000-0003-1999-6672), Kevin J. Anstrom, PhD (0000-0001-6452-2172),
Susan Halabi, PhD (0000-0003-4135-2777), John E. McKinnon, M.D., M.Sc (0000-0003-3916-
4021), Ravi Amaravadi, MD (0000-0002-5768-2474), Jorge Rojas Serrano, MD, PhD (0000-
0001-6980-7898), Benjamin S. Abella, MD MPhil (0000-0003-2521-0891), Angélica Margarita
Portillo-Vázquez, MD, MSc (0000-0002-5032-4076), Christopher W. Woods, MD, MPH,
Adrian Hernandez, MD, MHS (0000-0003-3387-9616), David R Boulware, MD MPH (0000-
0002-4715-0060), Susanna Naggie, MD, MHS (0000-0001-7721-6975), Radha Rajasingham,
MD (0000-0001-5531-0231)

Hwanhee Hong, Assistant Professor, Department of Biostatistics and Bioinformatics, Duke
University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA

Anne Friedland, Assistant Professor, Department of Infectious Disease, University of North
Carolina School of Medicine, 130 Mason Farm Rd, Chapel Hill, NC, 27514, USA

Mengyi Hu, Statistician, Department of Biostatistics and Bioinformatics, Duke University, 2424
Erwin Road, Ste 1105, Durham, NC, 27705, USA

Kevin J. Anstrom, Director, Collaborative Studies Coordinating Center, Gillings School of
Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA

Susan Halabi, Professor, Department of Biostatistics and Bioinformatics, Duke University, 2424
Erwin Road, Suite 11088, Durham, NC, 27705, USA

1
2
3 23 John E. McKinnon, Senior Staff, Infectious Diseases, Henry Ford Health, Clara Ford Pavilion,
4
5 24 Suite 322, 2799 W. Grand Blvd., Detroit, MI 48202, USA
6
7 25 Ravi Amaravadi, Professor of Medicine, University of Pennsylvania , 852 BRB 2/3 421 Curie
8
9 26 Blvd., Philadelphia, PA 19063, USA
10
11
12 27 Jorge Rojas Serrano, Attending physician, Interstitial Lung Disease and Rheumatology Unit,
13
14 28 Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas, Calzada de Tlalpan
15
16 29 4502, Tlalpan, Ciudad de México, México. 14080
17
18
19 30 Benjamin S. Abella, Professor, Department of Emergency Medicine, University of Pennsylvania,
20
21 31 423 Guardian Drive Room 412, Philadelphia, PA 19104, USA
22
23 32 Angélica Margarita Portillo-Vázquez, Attending physician, Otolaryngology department, Instituto
24
25 33 Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas. Calzada de Tlalpan 4502,
26
27 34 Tlalpan, Ciudad de México, México. 14080
28
29
30 35 Christopher W. Woods, Professor of Medicine, Duke University, 310 Trent Drive, Durham, NC
31
32 36 27710, USA
33
34
35 37 Adrian Hernandez, Professor of Medicine, Duke Clinical Research Institute, Duke University
36
37 38 300 W. Morgan Street Durham, NC 27701, USA
38
39
40 39 David R Boulware, Professor, Division of Infectious Diseases & International Medicine,
41
42 40 Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis, MN, 55455,
43
44 41 USA
45
46
47 42 Susanna Naggie, Professor of Medicine, Duke Clinical Research Institute, 300 West Morgan
48
49 43 Street, 5th Floor, Durham, NC 27710, USA
50
51
52
53
54
55
56
57
58
59
60

44 Radha Rajasingham, Assistant Professor, Division of Infectious Diseases & International
45 Medicine, Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis,
46 MN, 55455, USA

47
48 **Corresponding author:** Hwanhee Hong, Assistant Professor, Department of Biostatistics and
49 Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA,
50 hwanhee.hong@duke.edu

51
52 Abstract word count: 226

53 Word count: 2483

54
55 **Key words:** SARS-CoV-2, COVID-19, hydroxychloroquine, prophylaxis, health care workers,
56 meta-analysis, clinical trials

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective: We studied the safety and efficacy of hydroxychloroquine (HCQ) as pre-exposure prophylaxis for COVID-19 in healthcare workers (HCWs), using a meta-analysis of randomized controlled trials.

Data Sources: PubMed, and EMBASE databases were searched to identify randomized trials studying HCQ.

Study Selection: Ten randomized controlled trials (RCTs) were identified (n=5,079 participants).

Data Extraction and Synthesis: The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis between HCQ and placebo using a Bayesian random-effects model. A *pre-hoc* statistical analysis plan was written, and the review protocol was registered at PROSPERO (CRD42021285093)

Main Outcomes: The primary efficacy outcome was polymerase chain reaction (PCR)-confirmed SARS-CoV-2 infection and the primary safety outcome was incidence of adverse events. The secondary outcome included clinically suspected SARS-CoV-2 infection.

Results: Compared with placebo, HCWs randomized to hydroxychloroquine (HCQ) had no significant difference in PCR-confirmed SARS-CoV-2 infection (odds ratio [OR] 0.92, 95% credible interval [CI]: 0.58, 1.37) or clinically suspected SARS-CoV-2 infection (OR 0.78, 95% CI: 0.57, 1.10), and marginally significant difference in adverse events (OR 1.35, 95% CI: 1.03, 1.73).

Conclusions and Relevance: Our meta-analysis of ten RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in HCWs found that compared with placebo HCQ

does not significantly reduce the risk of confirmed or clinically suspected SARS-CoV-2 infection, while HCQ significantly increases adverse events.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This meta-analysis studied the safety and efficacy of hydroxychloroquine as pre-exposure prophylaxis for COVID-19 in healthcare workers.
- Bayesian meta-analysis models with random effects fitted the data.
- The ten trials included in the meta-analysis represent wide geographical locations including US, Canada, Mexico, India, Spain, Bolivia, Venezuela, Peru, and Pakistan.
- The findings can be applied to healthcare workers but should not be generalized to a broader population.

INTRODUCTION

Early during the SARS-CoV-2 pandemic, based on *in vitro* antiviral activity of both chloroquine and hydroxychloroquine against SARS-CoV-2 [1-3], clinicians considered use of hydroxychloroquine (HCQ) for treatment and prevention of SARS-CoV-2 infection and the associated disease, COVID-19. While there are now published randomized controlled trials of HCQ for the treatment of COVID-19 in the inpatient and outpatient setting [4, 5], there remains a lack of adequately powered randomized controlled trials of HCQ for the pre-exposure prophylaxis (PrEP) of SARS-CoV-2 infection. A number of COVID-19 clinical studies including PrEP studies were planned early in the pandemic; however, several never opened to enrollment and those that did open were closed early without reaching full accrual due to the rapidly

changing landscape of preventative therapies, including vaccines, and a significant shift in public opinion of HCQ as a medical intervention for SARS-CoV-2 [6].

Vaccination access remains insufficient globally [7]. Specifically, in low-income countries only 33% of healthcare workers are fully vaccinated. While high-income countries have better coverage, overall 38% of countries did not achieve the milestone of 70% vaccination coverage for healthcare workers by the end of 2021[8]. Thus, studying the pre-exposure prophylaxis potential for a drug with a known safety profile is crucial to protect people at high risk of exposures, such as healthcare workers (HCWs) [9, 10]. Two large randomized, placebo-controlled trials testing the safety and efficacy of HCQ as pre-exposure prophylaxis for COVID-19 in HCWs [11] [12], showed potential for a modest benefit of HCQ but were both underpowered, if a modest effect exists. More trials [13-15] studying HCQ as pre-exposure prophylaxis of COVID-19 in HCWs have been published with similar limitations.

To address the most common limitation, inadequate power to show a modest effect, we conducted a formal meta-analysis of pre-exposure prophylactic HCQ studies in HCWs. We conducted a systematic search for clinical trials of pre-exposure prophylactic use of HCQ against infection of SARS-CoV-2 in HCWs, thoroughly compared similarities and differences in characteristics of the identified studies and performed a Bayesian meta-analysis to combine results of the trials.

METHODS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis[16]. A statistical analysis plan was written in advance and the review protocol was registered at PROSPERO (CRD42021285093).

134

135 Search strategy and information sources

136 We searched PubMed/Medline and Ovid/Embase databases from database inception through the
137 final search date March 14, 2023. We used keywords related to COVID-19, HCQ, and
138 randomized controlled trials. The full search strategies are provided in eTable 1.

139

140 Eligibility criteria and study selection

141 The eligibility criteria included phase II or phase III randomized controlled trials (RCTs) of
142 hydroxychloroquine for use as pre-exposure prophylaxis in HCWs with moderate to high risk of
143 exposure. We excluded observational studies, crossover trials, studies where the method of
144 allocation to treatment was not truly random, duplicate studies, and non-original data studies. No
145 language, publication date, or publication status restrictions were applied. References of prior
146 systematic reviews and meta-analyses were also screened for related studies. Study selection
147 involved screening of titles and abstracts followed by full-text evaluation of possible eligible
148 studies.

149

150 Data collection process

151 Each of the selected studies were independently reviewed by two reviewers (AF, MH, or HH).
152 We extracted data on the study design, baseline characteristics, interventions, and outcomes. Any
153 disagreements of collected information between reviews were reconciled through discussion by
154 all three reviewers.

155

156 Outcome measures

The primary efficacy outcome for the meta-analysis was laboratory confirmed SARS-CoV-2 infection by polymerase chain reaction (PCR) test and the primary safety outcome was incidence of adverse events (Table 1). The secondary efficacy outcome was suspected or probable SARS-CoV-2 infection. Included studies had the following outcome definitions: (1) laboratory confirmed SARS-CoV-2 infection defined as COVID-19 like symptoms and positive SARS-CoV-2 PCR and (2) suspected or probable SARS-CoV-2 infection defined as COVID-19 like symptoms but lack of confirmatory PCR testing.

Table 1. Treatment strategies, adherence, trial-defined primary outcome, and study duration for trials included in the meta-analysis

| | Trial-defined primary outcome | Study duration | Treatment group | Randomized treatment assignment | Randomized sample size |
|--|--|---------------------|------------------|---|------------------------|
| Naggie et al.[13] (HERO-HCQ) | Confirmed (by NP swab PCR) or suspected COVID-19 infection through 30 days | 60 days | HCQ | HCQ 600 mg BID loading dose for Day 1, followed by 400 mg QD for 29 days | 683 |
| | | | Control | Placebo | 676 |
| Abella et al.[11] (PATCH) | COVID-19 infection as determined by positive NP swab over 8 weeks | 56 days (8 weeks) | HCQ | HCQ 600mg daily for 60 days | 64 |
| | | | Control | Placebo | 61 |
| Rajasingham et al.[12] (MN-COVID-PREP) | COVID-19 free survival time by lab confirmed or probable illness | 84 days (12 weeks) | HCQ ^a | HCQ loading doses (400 mg twice 6-8hrs apart), followed by 400 mg once weekly or 400 mg twice weekly for 84 days | 989 |
| | | | Control | Placebo | 494 |
| Rojas-Serrano et al.[14] | Time to symptomatic respiratory infection with a positive COVID RT PCR over 60 days | 60 days | HCQ | HCQ 200 mg daily for 60 days | 62 |
| | | | Control | Placebo | 65 |
| McKinnon et al.[15] (WHIP) | Lab confirmed cases of COVID-19 determined by either IgM and IgG serology in blood sample or RT-PCR test results | 56 days (8 weeks) | HCQ ^a | HCQ 400 mg loading dose for Day 1, followed by 200 mg daily or 400 mg weekly on the same day of each week for 56 days | 387 |
| | Confirmed new cases of COVID-19 | | Control | Placebo | 191 |
| Vijayaraghavan et al.[17] | Lab confirmed SARS-CoV-2 infection by PCR or presence of antibodies | 180 days (6 months) | HCQ | HCQ 400 mg twice on the day of enrollment, followed by 400 mg once a week for a total of 12 weeks plus personal | 213 |

| | | | | | |
|---------------------------|--|---------------------|------------------|---|-----|
| | | | | protective equipment (PPE) | |
| | | | Control | PPE | 203 |
| Polo et al.[18] (EPICOS) | Lab confirmed symptomatic COVID-19 by PCR | 84 days (12 weeks) | HCQ ^b | HCQ 200 mg once daily | 231 |
| | | | Control | Placebo | 223 |
| Llanos-Cuentas et al.[19] | COVID-19 cases confirmed by PCR or serological test | 28 days (4 weeks) | HCQ | HCQ loading dose of 600 mg on the first day, followed by 400 mg every other day plus PPE | 36 |
| | | | Control | PPE | 32 |
| Grau-Pujol et al.[20] | COVID-19 confirmed cases with seroconversion or PCR test | 180 days (6 months) | HCQ | HCQ 400 mg daily for the four consecutive days, followed by 400 mg weekly | 142 |
| | | | Control | Placebo | 127 |
| Syed et al.[17] | COVID-19-free survival (COVID-19 confirmed by PCR) | 84 days (12 weeks) | HCQ ^a | HCQ 400 mg twice for Day 1, followed by 400 mg weekly or HCQ 400 mg once every 3 weeks or HCQ 200 mg once every 3 weeks | 154 |
| | | | Control | Placebo | 46 |

HCQ=Hydroxychloroquine

^a More than one HCQ groups with different doses are lumped.

^b The Polo et al. study randomized participants to four treatment groups, and the HCQ and control groups are used in our meta-analysis.

Treatment assignment

Our meta-analysis did not study HCQ dosing specific effects. For studies randomizing participants to more than one HCQ arm with different doses, all HCQ arms were merged and considered as a single HCQ arm. Such studies include the Rajasingham et al., McKinnon et al. and Syed et al. studies.

Risk of bias and certainty of evidence assessment

Two independent reviewers (AF, HH) assessed the risk of bias (low, intermediate, high) of the included studies using the Cochrane's Collaboration tool [21] (eTable 2). We assessed the certainty of evidence using the Grades of Recommendation Assessment, Development and Evaluation (GRADE) approach [22].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Statistical analysis

Bayesian logistic regression meta-analysis models under two assumptions (fixed effect and random effects) were fitted to estimate the odds ratio of having an outcome between hydroxychloroquine and placebo [23]. The fixed effect model assumes that the odds ratio is constant across studies, while the random effects model accounts for heterogeneity in the odds ratios across studies. To assess and compare the goodness-of-fit of the fitted fixed and random effects models, we calculated the Watanabe-Akaike information criterion [24]. In the Bayesian models, we assigned non-informative prior distributions as no prior information was available. The odds ratios and the associated 95% credible intervals were estimated using Markov chain Monte Carlo (MCMC) algorithms. In addition, we calculated Bayesian posterior probabilities of the odds ratio smaller than 1 or 0.5 for the primary efficacy outcome, and greater than 2 for the safety outcome [25]. The standard deviation of the random effects and I^2 [26] were estimated to quantify the between-study heterogeneity, where small values of both metrics indicate slight heterogeneity. To identify publication bias, we plotted and assessed funnel plots for their symmetry, and conducted the Egger's test[27]. All Bayesian meta-analyses were conducted using the `rstan` package (version 2.21.2)[28] in R 4.0.2 [29]. We used two parallel chains, where each chain consists of 50,000 samples after a 25,000-sample burn-in. We checked convergence of the MCMC chains for all model parameters using trace plots and Gelman-Rubin diagnostic statistics [30].

Patient and public involvement

No patient involved.

208 RESULTS

209 Search results

210 Our database search resulted in 350 unique studies after excluding duplicates. Of those, 339
211 studies were screened out due to irrelevance based on title and abstract screening. Eleven studies
212 were assessed in full-text for eligibility (Figure 1). Of those, one trial was excluded from the
213 meta-analysis because it studied with non-healthcare worker populations. As a result, a total of
214 ten studies in a population consisting of HCWs were identified (Table 1).

215

216 Study and patient characteristics

217 Study design, population, treatment strategies, and key characteristics are presented in Table 1
218 and eTable 3. A total of 5,079 randomized participants (2,961 randomized to HCQ) from the 10
219 studies were included in the meta-analysis. The ten studies defined HCWs broadly and included
220 first responders (emergency medical services, fire, and police). The follow-up duration of the 10
221 studies ranged from 28 days to 180 days. The HCQ dosing scheme varied across studies,
222 including daily dosing ranging from 200 to 600mg daily with or without a loading dose and once
223 or twice weekly or once every three weeks dosing. The duration of therapy also varied across
224 studies (Table 1). The trial-specific definitions of primary outcome and adverse events are
225 comparable across trials (Table 1, eTable 4).

226

227 Baseline characteristics by randomized treatment assignment are reported (eTable 5). The
228 average age ranged between 31 and 45. The aggregate proportion of women within each study
229 varied across the 10 trials, with a range from 44% to 69%. In addition, the Abella et al. and
230 Rojas-Serrano et al. studies had smaller sample size compared with the other three studies and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

showed a difference in female ratio between placebo and HCQ groups. In the Naggie et al.,
Abella et al., Rajasingham et al., and McKinnon et al., studies, over 80% of study participants
were white. The Abella et al. and Rajasingham et al. studies had high proportions of HCWs
working in an emergency department (56% and 41%, respectively) and the Abella et al. study
had a high proportion of nurses (67%).

Several studies reported treatment adherence assessed by two methods: self-reported adherence
and/or pill count at the end of the study. The Rajasingham et al. study additionally conducted
remote blood sampling to verify HCQ concentrations in a subset. Adherence varied significantly
across the studies, with a low proportion of approximately 52% in the Rojas-Serrano et al. study
and 97-98% in the Abella et al. study.

Results of meta-analysis

Overall, 3.4% (171/5039) developed PCR-confirmed SARS-CoV-2 infection and 5.6% (230/4087)
developed suspected COVID-19 that was not laboratory confirmed. Since the goodness-of-fit
assessment using Watanabe-Akaike information criterion concluded that the random effects meta-
analysis model was as good as or better than the fixed effect meta-analysis model for all outcomes,
we reported the results under the random effects model. Compared with placebo, HCWs
randomized to HCQ had numerically lower rate of PCR-confirmed SARS-CoV-2 infection cases
(odds ratio [OR] 0.92, 95% credible interval [CI]: 0.58, 1.37), and suspected or probable SARS-
CoV-2 infection cases (OR 0.78, 95% CI: 0.57, 1.10). None of these odds ratios were statistically
significant. Participants treated with HCQ had a numerically higher rate of adverse events (OR
1.35, 95% CI: 1.03, 1.73) with marginally statistical significance (Figure 2). The outcome data

used in our analyses are presented in eTable 6. The GRADE scores for the odds ratios with respect to all three outcomes were downgraded by 1 due to wide credible intervals of odds ratios, resulting in moderate certainty of evidence.

The Bayesian posterior probabilities of the odds ratio less than 1 for the confirmed SARS-CoV-2 infection outcome (i.e., the probability of HCQ favoring over placebo) was 0.67, while the posterior probability of odds ratio less than 0.5 (i.e., the probability that the odds of having a confirmed SARS-CoV-2 infection outcome in HCQ is less than a half of the odds in placebo) was 0.009. The posterior probability of the odds ratio greater than 2 for the adverse event outcome (i.e., the probability that the odds of having an adverse event in HCQ is greater than twice of the odds in placebo) was 0.004.

Our meta-analysis showed little or moderate variability of effect estimates across studies with I^2 value of 0%, 0%, and 43%, and the estimated standard deviation of the random effects of 0.39, 0.26, and 0.45 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively. Funnel plots (eFigure) showed no indication of publication bias and the associated Egger's test results supported that the funnel plots were not asymmetry with p-values of 0.308, 0.305, and 0.794 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively.

DISCUSSION

Understanding the pre-exposure prophylactic effect of HCQ against COVID-19 remains relevant, as its use continues, particularly in the international setting [31, 32]. Our meta-analysis

1
2
3 277 of the ten RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in
4
5 278 5,079 HCWs found that HCQ did not have a statistical association with fewer confirmed or
6
7
8 279 suspected/probable SARS-CoV-2 infection cases compared with placebo. The geographical
9
10 280 locations of the 10 trials included in the meta-analysis are US, Canada, Mexico, India, Spain,
11
12 281 Bolivia, Venezuela, Peru, and Pakistan (eTable 3). While the odds ratios of most studies favor
13
14 282 HCQ, the credible intervals remain wide suggesting low certainty in the true point estimate. Two
15
16 283 studies including the Llanos-Cuentas et al. study conducted in Peru and the Syed et al. study
17
18 284 conducted in Pakistan showed odds ratios favoring placebo, though the credible intervals remain
19
20 285 wide. Furthermore, in this population, COVID-19 events rates were low, particularly for the
21
22 286 most relevant PCR-confirmed infection outcome. The low event rate raises further concern for
23
24 287 the uncertainty of these outcomes. Thus, if there is a minimal effect, the absolute benefit would
25
26 288 be low. To gain more certainty, a very large study would need to be done and this is difficult to
27
28
29 289 support now due to availability of highly effective vaccines. The safety profile of HCQ in the
30
31 290 outpatient setting is well understood [33]. In these outpatient studies there was marginally
32
33 291 statistically significant difference in adverse events in the HCQ versus the placebo arm,
34
35
36 292 indicating that HCQ is less safe than placebo.
37
38
39
40 293

41
42 294 Our findings can be applied to HCWs but should not be generalized to a broader population. Our
43
44 295 systematic search found only one published RCT of pre-exposure prophylaxis for non-healthcare
45
46 296 worker populations and the study were excluded from our meta-analysis. This study was
47
48 297 conducted in Singapore [34] and showed a significant reduction in the risk of COVID-19
49
50 298 infection in the HCQ arm when compared with the comparator arm, vitamin C. However, this
51
52 299 study showed moderate risk of bias as it used an open-label cluster-randomization design, the
53
54
55
56
57
58
59
60

Institutional Review Board excluded higher risk persons from the hydroxychloroquine arm only, and the participants may not be representative of a general population due to the communal living environment.

A Bayesian meta-analysis approach was used to fit the data. The Bayesian meta-analysis approach has several advantages. First, its flexibility and the MCMC sampling methods to estimate posterior distributions provide probability-based quantities (e.g., posterior probability of an odds ratio smaller than 0.5) that complement typical meta-analysis results (e.g., odds ratios and the associated credible intervals) and help decision making [35]. Second, the Bayesian meta-analysis model with random effects estimates the between-study variability better than the frequentist counterparts [36]. Third, when it comes to with binary outcomes, the Bayesian approach handles rare events better than the frequentist counterparts [23].

A recently published meta-analysis by García-Albéniz et al. [37] investigated pre-exposure (seven RCTs included) and post-exposure (four RCTs included) prophylactic effects of HCQ, but not limited to the HCW population. They found significant pre-exposure prophylactic effects of HCQ on SARS-CoV-2 infection, different from ours. The seven pre-exposure prophylaxis RCTs included in the García-Albéniz et al. meta-analysis consisted of six RCTs that were in our meta-analysis and the aforementioned Singapore study that was excluded from our meta-analysis. Our meta-analysis provides the most up-to-date, systematic, and comprehensive evidence about prophylactic effects of HCQ focusing on the HCW population.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

322 Although a meta-analysis allows for combining evidence from multiple studies in a principled
323 way, our meta-analysis has limitations. First, our analysis did not evaluate effects of different
324 HCQ doses and combined multiple HCQ arms using different doses in three studies. The RCTs
325 included in our meta-analysis studied varying dosing schemes and a meta-analysis using
326 aggregate-level data is not a sufficient source to study dosing effects. Second, detailed subgroup
327 analyses were not conducted due to limited information. Individual-level data are required to
328 study both dosing and subgroup effects.

329
330 Our meta-analysis of ten RCTs investigating safety and efficacy of HCQ as pre-exposure
331 prophylaxis in HCWs provides the most up-to-date evidence on HCQ. Although most individual
332 trials were underpowered and showed null data, integrating the results systematically via meta-
333 analysis contributes to the scientific literature and provides certain answers to the question. We
334 found that HCQ does not reduce the risk of confirmed or probable SARS-CoV-2 infection, but
335 increase risk of adverse events compared with placebo. Hydroxychloroquine should not be used
336 for pre-exposure prophylaxis in the HCW population.

337
338 **Contributors**
339 All authors fulfill the ICMJE criteria for authorship. HH, SN, RR, and KJA designed the study.
340 HH, AF, and MH collected and analyzed the data. HH, SN, and RR wrote the manuscript. SH
341 and KJA provided statistical review and AF, JEM, RA, JRS, BSA, AMPV, CWW, AH and DRB
342 provided clinical review. All authors approved and decided to submit the paper for publication.
343 Hwanhee Hong – HH
344 Anne Friedland – AF

345 Mengyi Hu – MH

346 Kevin J. Anstrom – KJA

347 Susan Halabi – SH

348 John E. McKinnon – JEM

349 Ravi Amaravadi – RA

350 Jorge Rojas Serrano – JRS

351 Benjamin S. Abella – BSA

352 Angélica Margarita Portillo-Vázquez – AMPV

353 Christopher W. Woods – CWW

354 Adrian Hernandez – AH

355 David R Boulware – DRB

356 Susanna Naggie – SN

357 Radha Rajasingham – RR

358 **Funding**

359 This study is funded by the Patient Centered Outcomes Research Institute (PCORI), Contract
360 Number COVID-19-2020-001. The funder had no role in the design, conduct, analysis, or
361 reporting of this study.

362 **Competing interests**

363 All authors except Dr. Abella reported no financial relationship with commercial interest. Dr.
364 Abella have received NIH funds for COVID-19 related research, and holds equity in VOC
365 Health, a start-up company that is developing novel covid testing.

366 **Ethics Approval**

Ethics approval was not required because this study used publicly available aggregate data that were not involved with patients’ information or prospective data collection.

Data sharing statement

The data are presented in eTable 6.

REFERENCES

1. Kalil, A.C., *Treating COVID-19—off-label drug use, compassionate use, and randomized clinical trials during pandemics*. JAMA, 2020. **323**(19): p. 1897-1898.
2. McCreary, E.K., J.M. Pogue, and o.b.o.t.S.o.I.D. Pharmacists, *Coronavirus Disease 2019 Treatment: A Review of Early and Emerging Options*. Open Forum Infectious Diseases, 2020. **7**(4).
3. Wang, M., et al., *Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro*. Cell research, 2020. **30**(3): p. 269-271.
4. RECOVERY Collaborative Group, *Effect of hydroxychloroquine in hospitalized patients with Covid-19*. New England Journal of Medicine, 2020. **383**(21): p. 2030-2040.
5. Skipper, C.P., et al., *Hydroxychloroquine in nonhospitalized adults with early COVID-19: a randomized trial*. Annals of internal medicine, 2020. **173**(8): p. 623-631.
6. Halabi, S., et al., *Landscape of coronavirus disease 2019 clinical trials: New frontiers and challenges*. Clinical Trials, 2022: p. 17407745221105106.
7. Padma, T., *COVID vaccines to reach poorest countries in 2023—despite recent pledges*. Nature, 2021. **595**(7867): p. 342-343.
8. Nabaggala, M.S., et al., *The global inequity in COVID-19 vaccination coverage among health and care workers*. International Journal for Equity in Health, 2022. **21**(3): p. 147.
9. World Health Organization. *Prevention, identification and management of health worker infection in the context of COVID-19*. 2020 [cited 2022 May 13th]; Available from: <https://www.who.int/publications/i/item/10665-336265>
10. The United Kingdom Office for National Statistics. *Coronavirus (COVID-19) infections in the community in England: May 2021*. 2021 [cited 2022 May 13th]; Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristicsofpeopletestingpositiveforcovid19incountriesoftheuk20may2021#percentage-testing-positive-for-covid-19-by-patient-facing-and-non-patient-facing-job-roles-uk>.
11. Abella, B.S., et al., *Efficacy and safety of hydroxychloroquine vs placebo for pre-exposure SARS-CoV-2 prophylaxis among health care workers: a randomized clinical trial*. JAMA internal medicine, 2021. **181**(2): p. 195-202.
12. Rajasingham, R., et al., *Hydroxychloroquine as Pre-exposure Prophylaxis for Coronavirus Disease 2019 (COVID-19) in Healthcare Workers: A Randomized Trial*. Clinical Infectious Diseases, 2020. **72**(11): p. e835-e843.

13. Naggie, S., et al., *Hydroxychloroquine for pre-exposure prophylaxis of COVID-19 in health care workers: a randomized, multicenter, placebo-controlled trial* *Healthcare Worker Exposure Response and Outcomes of Hydroxychloroquine (HERO-HCQ)*. International Journal of Infectious Diseases, 2023. **129**: p. 40-48.
14. Rojas-Serrano, J., et al., *Hydroxychloroquine for prophylaxis of COVID-19 in health workers: A randomized clinical trial*. PLoS One, 2022. **17**(2): p. e0261980.
15. McKinnon, J.E., et al., *Safety and tolerability of hydroxychloroquine in health care workers and first responders for the prevention of COVID-19: WHIP COVID-19 Study*. International Journal of Infectious Diseases, 2022. **116**: p. 167-173.
16. Hutton, B., et al., *The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations*. Annals of internal medicine, 2015. **162**(11): p. 777-784.
17. Tirupakuzhi Vijayaraghavan, B.K., et al., *Hydroxychloroquine plus personal protective equipment versus personal protective equipment alone for the prevention of laboratory-confirmed COVID-19 infections among healthcare workers: a multicentre, parallel-group randomised controlled trial from India*. BMJ Open, 2022. **12**(6): p. e059540.
18. Polo, R., et al., *Daily tenofovir disoproxil fumarate/emtricitabine and hydroxychloroquine for pre-exposure prophylaxis of COVID-19: a double-blind placebo-controlled randomized trial in healthcare workers*. Clinical Microbiology and Infection, 2023. **29**(1): p. 85-93.
19. Llanos-Cuentas, A., et al., *Hydroxychloroquine to prevent SARS-CoV-2 infection among healthcare workers: early termination of a phase 3, randomised, open-label, controlled clinical trial*. BMC Research Notes, 2023. **16**(1): p. 22.
20. Grau-Pujol, B., et al., *Pre-exposure prophylaxis with hydroxychloroquine for COVID-19: a double-blind, placebo-controlled randomized clinical trial*. Trials, 2021. **22**(1): p. 808.
21. Sterne, J.A., et al., *RoB 2: a revised tool for assessing risk of bias in randomised trials*. bmj, 2019. **366**.
22. Puhan, M.A., et al., *A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis*. Bmj, 2014. **349**.
23. Hong, H., C. Wang, and G.L. Rosner, *Meta-analysis of rare adverse events in randomized clinical trials: Bayesian and frequentist methods*. Clinical Trials, 2021. **18**(1): p. 3-16.
24. Watanabe, S. and M. Oppel, *Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory*. Journal of machine learning research, 2010. **11**(12).
25. Ferreira, D., et al., *Bayesian predictive probabilities: a good way to monitor clinical trials*. British journal of anaesthesia, 2021. **126**(2): p. 550-555.
26. Higgins, J.P. and S.G. Thompson, *Quantifying heterogeneity in a meta-analysis*. Statistics in medicine, 2002. **21**(11): p. 1539-1558.
27. Egger, M., et al., *Bias in meta-analysis detected by a simple, graphical test*. Bmj, 1997. **315**(7109): p. 629-634.
28. Stan Development Team, *RStan: the R interface to Stan*. R package version, 2020. **2.21.2**.
29. R Core Team, *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2021.
30. Gelman, A. and D.B. Rubin, *Inference from iterative simulation using multiple sequences*. Statistical science, 1992. **7**(4): p. 457-472.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

449 31. Infante, M., et al., *Hydroxychloroquine in the COVID-19 pandemic era: in pursuit of a*
450 *rational use for prophylaxis of SARS-CoV-2 infection*. Expert review of anti-infective
451 therapy, 2021. **19**(1): p. 5-16.

452 32. *Revised advisory on the use of hydroxychloroquine (HCQ) as prophylaxis for SARS-CoV-2*
453 *infection (in supersession of previous advisory dated 23rd March. 2020)*. 2022; Available
454 from:
455 [https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ](https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ_SARS_CoV2_infection.pdf)
456 [SARS CoV2 infection.pdf](https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ_SARS_CoV2_infection.pdf).

457 33. Lofgren, S.M., et al. *Safety of hydroxychloroquine among outpatient clinical trial*
458 *participants for COVID-19*. in *Open forum infectious diseases*. 2020. Oxford University
459 Press US.

460 34. Seet, R.C.S., et al., *Positive impact of oral hydroxychloroquine and povidone-iodine*
461 *throat spray for COVID-19 prophylaxis: An open-label randomized trial*. International
462 Journal of Infectious Diseases, 2021. **106**: p. 314-322.

463 35. Hong, H., et al., *A Bayesian missing data framework for generalized multiple outcome*
464 *mixed treatment comparisons*. Research synthesis methods, 2016. **7**(1): p. 6-22.

465 36. Hong, H., et al., *Comparing Bayesian and frequentist approaches for multiple outcome*
466 *mixed treatment comparisons*. Medical Decision Making, 2013. **33**(5): p. 702-714.

467 37. García-Albéniz, X., et al., *Systematic review and meta-analysis of randomized trials of*
468 *hydroxychloroquine for the prevention of COVID-19*. European Journal of Epidemiology,
469 2022. **37**(8): p. 789-796.

Figure Legends

Figure 1. Flowchart of literature review

Figure 2. Forest plots of the meta-analysis results showing the number of events (y), sample size (n), posterior median of odds ratios, and the associated 95% credible intervals comparing HCQ versus placebo for (a) lab-confirmed positive COVID-19, (b) suspected COVID-19, and (c) adverse events.

For peer review only

1 Identification
2
3
4
5

6
7
8
9
10
11


12 Screening
13
14
15
16
17
18


19
20
21
22
23
24
25

26 Eligibility
27
28
29
30
31
32
33


34
35
36
37
38
39

40 Included
41
42
43
44
45
46
47
48






(b) Suspected COVID-19



(c) Adverse events



Supplementary Materials

CONTENTS

- eTable 1. Search code
- eTable 2. Risk of bias
- eTable 3. Characteristics of included trials
- eTable 4. Definition of adverse events
- eTable 5. Baseline characteristics
- eTable 6. Results of outcome measures in each study
- eFigure. Funnel plots for the three outcomes



















































eTable 1. Search code that was used to identify publications as of March 14, 2023**PubMed search**

| | |
|----|--|
| #1 | covid[Title] OR coronavirus[Title] OR sars-cov-2[Title] |
| #2 | hydroxychloroquine[Title] |
| #3 | randomized[Title/Abstract] OR randomized[Title/Abstract] |
| #4 | #1 AND #2 AND #3 |

Embase search

| | |
|----|---|
| #1 | covid:ti OR coronavirus:ti OR 'sars cov 2':ti |
| #2 | hydroxychloroquine:ti |
| #3 | randomized:ab,ti OR randomised:ab,ti |
| #4 | #1 AND #2 AND #3 |

eTable 2. Risk of bias for trials included in the meta-analysis using the Cochrane risk assessment tool. Green circle is for low risk and yellow circle is for some concerns

| | Selection bias (Randomization process) | Performance bias (Deviations from the intended interventions) | Attrition bias ¹ (Missing outcome data) | Reporting bias (Measurement of the outcome) | Other sources of bias (Selection of the reported result) |
|---------------------------------------|---|---|---|---|---|
| Naggie et al. (HERO-HCQ) |  |  |  |  |  |
| Abella et al. (PATCH) |  |  |  |  |  |
| Rajasingham et al. (MN-COVID-PREP) |  |  |  |  |  |
| Rojas-Serrano et al. |  |  |  |  |  |
| McKinnon et al. (WHIP) |  |  |  |  |  |
| Vijayaraghavan et al. |  |  |  |  |  |
| Polo et al. (EPICOS) |  |  |  |  |  |
| Llanos-Cuentas et al. |  |  |  |  |  |
| Grau-Pujol et al. |  |  |  |  |  |
| Syed et al. |  |  |  |  |  |

¹ The Rojas-Serrano et al. study reported minimal loss to follow-up (<10%). The Rojas-Serrano et al. study reported 18% (25/130) lost to follow-up and additional 12% (16/130) discontinued the intervention.

eTable 3. Characteristics of trials included in the meta-analysis

| | Naggie et al. (HERO-HCQ) | Abella et al. (PATCH) | Rajasingham et al. (MN-COVID-PREP) | Rojas-Serrano et al. | McKinnon et al. (WHIP) |
|---|--|--|--|---|--|
| N (randomization) | 1360 | 132 | 1496 | 130 | 624 |
| Study start date ¹ | 4/22/2020 | 4/9/2020 | 4/6/2020 | 4/21/2020 | 4/10/2020 |
| Study completion date ² | 1/9/2021 | 11/13/2020 | 7/13/2020 | 3/31/2021 | 12/14/2020 |
| Occupation | HCWs at risk of COVID exposure through work in the ICU, emergency department, emergency services, respiratory services or COVID unit | HCWs (Physicians, nurses, certified nursing assistants, emergency technicians, respiratory therapists) eligible working >20 hrs/week | HCWs (physicians, nurses, emergency medical technicians) with direct contact with COVID patients including emergency department and ICU setting, first responders and performing aerosol generating procedures | HCWs (nurses, nursing aids, cleaning staff, orderlies, respiratory therapists and physicians) taking care of hospitalized patients with COVID | HCW, first responders and correlational/law officers, nursing home workers, medical students, public transit workers, household family members of HCW in Michigan and Ohio |
| Sites | 34 sites across the US | 2 tertiary urban hospitals | Multiple sites nationwide across US and Canada | Single site (National Institute of Respiratory Diseases of Mexico) | Multiple sites at Michigan in the US |
| Randomization | Yes (Phase III) | Yes (Phase II) | Yes (Phase III) | Yes (Phase III) | Yes (Phase III) |
| Trial type | Double-blinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | No weight requirement | <40kg excluded | <50kg excluded | N/A |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Excluded | Excluded | Excluded | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Included | Exclude |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Included | |
| Significant renal disease (stage IV or greater) | Excluded | Included | Excluded | Excluded | Exclude |
| Pregnant/breastfeeding | Included | Excluded | Included in US, Excluded in Canada | Excluded | Exclude |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Excluded | Included | Exclude |
| Use of other medications with significant drug interactions | Included | Excluded | Excluded | Included | N/A |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine or azithromycin) | Any treatment for COVID-19 within 14 days excluded | Current use of HCQ or chloroquine excluded | HCQ or chloroquine within 30 days excluded | Chronic use of HCQ included |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | N/A | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Excluded | Excluded if symptoms within 2 weeks unless negative COVID test | Excluded | Excluded | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | N/A |
| Positive COVID serology | Included | Included | N/A | Included | N/A |
| Analysis | Modified intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

| | Vijayaraghavan et al. | Polo et al. (EPICOS) | Llanos-Cuentas et al. | Grau-Pujol et al. | Syed et al. |
|---|---|--|--|--|--|
| N (randomization) | 416 | 454 | 68 | 269 | 200 |
| Study start date ¹ | 6/29/2020 | 4/2020 Spain, 10/2020 Bolivia, 3/2021 Venezuela | June, 2020 | 4/4/2020 | 5/1/2020 |
| Study completion date ² | 2/4/2021 | 5/30/2021 | November, 2020 | Study halted a 1 month analysis | Not reported |
| Occupation | HCWs in an environment with exposure to COVID-19 (physicians, nurses, allied health workers and ancillary health workers) | HCWs (physicians, nurses, medical students, other workers with and without direct patient contact) | HCWs (physicians, nursing staff, technical staff and nursing assistants involved in care of COVID-19 patients) | HCWs (physicians, nurses, nurse assistants and administrators working at least 3 days a week in the trial hospitals) | HCWs at risk of COVID-19 exposure including physicians, nurses, first responders, those performing aerosol generating procedures or working in the emergency department, ICU, and general medicine wards |
| Sites | 9 hospitals across India | Multiple sites across Spain, Venezuela and Bolivia | 4 public hospitals across the Lima metropolitan area | 3 hospitals in Barcelona, Spain | Single hospital in Pakistan |
| Randomization | Yes | Yes | Yes (Phase III) | Yes | Yes (Phase II) |
| Trial type | Unblinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18-70 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | <40kg excluded | No weight requirement | No weight requirement | <40 kg |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Included | Excluded | Included | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Excluded | Excluded |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Excluded | Excluded |
| Significant renal disease (stage IV or greater) | Included | Excluded | Excluded | Excluded | Excluded |
| Pregnant/breastfeeding | Excluded | Excluded | Included | Excluded | Excluded |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Included | Excluded | Excluded |
| Use of other medications with significant drug interactions | Excluded | Included | Included | Excluded | Excluded |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine azithromycin) | Any medication as prophylaxis against COVID-19 after 3/1/21 | Use of hydroxychloroquine, chloroquine or azithromycin in the last 30 days excluded | Treatment with chloroquine or hydroxychloroquine within the last 1 month | Those already taking hydroxychloroquine were excluded |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | Excluded | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Not specified in exclusion criteria | Excluded | Not specified in exclusion criteria | Not specified in exclusion criteria | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | Excluded |
| Positive COVID serology | N/A | N/A | N/A | Excluded | Excluded |
| Analysis | Intention-to-treat | Not reported | Intention-to-treat | Intention-to-treat | Not reported |

HCW=Healthcare workers; ICU=Intensive care unit; ¹ Date when first participant was enrolled; ² Date when final data were collected for the last participant

eTable 4. Definition of adverse events

| Trial | AE definition |
|---|---|
| Naggie et al. (HERO-HCQ) | Adverse events include general disorders and administration site conditions, psychiatric disorders, skin and subcutaneous tissue disorders, cardiac disorders, infections and infestations, nervous system disorders, gastrointestinal disorders, investigations (electrocardiogram QT prolonged and heart rate increased), ear and labyrinth disorders, renal and urinary disorders, and respiratory, thoracic and mediastinal disorders. |
| Abella et al. (PATCH) | Adverse events include abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. |
| Rajasingham et al. (MN-COVID-PREP) | Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and others. |
| Rojas-Serrano et al. | Examples of adverse events are as follows: abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and other. |
| McKinnon et al. (WHIP) | Covid-19 related symptoms, covid-19 clinical disease and medication adverse effects including gastrointestinal disorders, nervous system disorders, respiratory, thoracic and mediastinal disorders, general disorders and administration site conditions, cardiac disorders, musculoskeletal and connective tissue disorders, psychiatric disorders, skin and subcutaneous tissue disorders, ear and labyrinth disorders, and eye disorders. |
| Vijayaraghavan et al. | Adverse events listed in each category at the participant level were categorized as cardiac, gastro-intestinal, headache, and Qtc prolongation. |
| Polo et al. (EPICOS) | Adverse events were classified by organ system and included: gastrointestinal disorders, blood and lymphatic system disorders, cardiac disorders, ear and labyrinth disorders, eye disorder, general disorders, immune system disorder, infections, injuries, investigations, metabolism and nutrition disorders, musculoskeletal/connective tissue disorders, nervous system disorders, psychiatric disorders, renal and urinary disorders, reproductive system disorders, respiratory disorders, skin disorders and vascular disorders. |
| Llanos-Cuentas et al. | Adverse events from grade 1 to grade 3 and above. Note that the Llanos-Cuentas et al. study did report the number of adverse events (not participants) in the HCQ group only. Due to limited information, it was excluded from the meta-analysis with the adverse event outcome. |
| Grau-Pujol et al. | Adverse events included: general symptoms (fever, chills, sweating, malaise, myalgia, arthralgia), gastrointestinal symptoms (nausea, abdominal pain, diarrhea, dysgeusia), dermatological symptoms (itching, rash), respiratory symptoms (rhinorrhea, sore throat / odynophagia, cough, pleuritic pain, dyspnea), neurologic symptoms (headache, visual disturbances), and cardiovascular symptoms. Events were graded mild, moderate and severe. |
| Syed et al. | Syed et al. report the number of patients in each group who experienced adverse events, but did not report what the events were. Due to limited information, it was excluded from the meta-analysis with the adverse event outcome. |

eTable 5. Baseline characteristics with additional variables and detailed information. Sample mean and standard deviation (in parenthesis) are reported for continuous variables, and the number of participants and proportion (in parenthesis) are reported for binary or categorical variables.

| | | Naggie et al. (HERO-HCQ) | | Abella et al. (PATCH) | | Rajasingham et al. (MN-COVID-PREP) | | Rojas-Serrano et al. | | McKinnon et al. (WHIP) | |
|--------------------|---|-----------------------------|--------------------|--------------------------|-------------------------|---------------------------------------|--------------------------|-----------------------------|-------------------------------|--|-------------|
| | | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo |
| N (randomization) | | 683 | 676 | 66 | 66 | 989 | 494 | 62 | 65 | 387 | 191 |
| Age | | 44.2 (11.9) | 43.1 (11.2) | 31 (20-66) ³ | 34 (23-62) ³ | 41.5 (35, 49) ³ | 40 (34, 48) ³ | 31.0 (26.4-39) ⁴ | 31.9 (27.2-43.7) ⁴ | 45.7 (11.6); 44.9 (11.4) ² | 44.1 (12.7) |
| Female | | 442 (64.7%) | 446 (66.0%) | 54 (82%) | 37 (56%) | 519 (52.5%) | 241 (48.8%) | 29 (42.6%) | 42 (64.6%) | 220 (57%) | 114 (60%) |
| BMI (kg/m^2) | | 28.3 (6.3) | 28.6 (6.7) | 26 (19-37) ⁵ | 26 (20-50) ⁵ | | | 26.7 (3.9) | 27.2 (4.6) | | |
| Current smoker | | | | 0 (0%) | 0 (0%) | 38 (3.84%) | 13 (2.6%) | 20 (32.2%) ⁶ | 23 (35.4%) ⁶ | | |
| Race/ Ethnicity | White | 624 (91.4%) | 610 (90.2%) | 55 (83%) | 54 (82%) | 852 (86.1%) | 419 (84.8%) | | | 334 (86%) | 161 (84%) |
| | Asian | | | 7 (11%) | 7 (11%) | 46 (4.7%) | 29 (5.9%) | | | 23 (6%) | 15 (8%) |
| | African American | 18 (2.6%) | 23 (3.4%) | 3 (4%) | 1 (2%) | 10 (1.0%) | 10 (2.0%) | | | 15 (4%) | 9 (5%) |
| | Hispanic | 39 (5.7%) | 40 (5.9%) | 0 (0%) | 2 (3%) | 40 (4.0%) | 18 (3.6%) | | | 11 (3%) | 7 (4%) |
| Comorb idities | Asthma | 58 (8.5%) | 77 (11.4%) | 9 (14%) | 14 (21%) | 91 (9.2%) | 59 (11.9%) | | | | |
| | Diabetes | 20 (2.9%) | 35 (5.2%) | 1 (2%) | 3 (5%) | 36 (3.6%) | 14 (2.8%) | | | | |
| | Hypertension | 99 (14.5%) | 99 (14.6%) | 3 (5%) | 14 (21%) | 145 (14.7%) | 60 (12.1%) | | | | |
| | None | | | 54 (82%) | 40 (61%) | 646 (65.3%) | 336 (68.0%) | 53 (85.5%) | 58 (89.2%) | | |
| Practice Location | Emergency Department | 96 (14.1%) | 94 (13.9%) | 38 (58%) | 36 (55%) | 417 (42.2%) | 190 (38.5%) | | | 48 (12%) | 19 (10%) |
| | Internal Medicine ward | | | 17 (26%) | 18 (27%) | 98 (9.9%) | 56 (11.3%) | | | 31 (8%) | 20 (10%) |
| | ICU/anesthesia | | | 6 (9%) | 6 (9%) | | | | | | |
| | Labor and delivery | | | 5 (7%) | 6 (9%) | | | | | | |
| | Ambulance | 66 (9.7%) | 63 (9.3%) | | | 73 (7.4%) | 45 (9.1%) | | | | |
| | Congregate care setting | | | | | 46 (4.7%) | 20 (4.0%) | | | | |
| | ICU | 48 (7.0%) | 59 (8.7%) | | | 184 (18.6%) | 85 (17.2%) | | | 37 (10%) | 23 (12%) |
| | Operating room | | | | | 103 (10.4%) | 75 (15.2%) | | | | |
| | EMS, Fire and Police | | | | | | | | | 32 (8%) | 16 (8%) |
| | First Responders | | | | | | | | | | |
| Occupation | Nurse | 186/677 (27.5%) | 167/668 (25.0%) | 46 (70%) | 42 (64%) | | | | | | |
| | Physician | 143/677 (21.1%) | 144/668 (21.6%) | 11 (17%) | 16 (24%) | | | | | | |
| | Certified Nurse Assistant | | | 2 (3%) | 2 (3%) | | | | | | |
| | ED Technician | | | 3 (4%) | 1 (2%) | | | | | | |
| | Respiratory therapist | 15/677 (2.2%) | 18/668 (2.7%) | 3 (4%) | 5 (7%) | | | | | | |
| | Nurse or Physician | | | | | | | 31 (50%) | 33 (50.8%) | | |
| | Emergency Medicine Provider | | | | | 407 (41.1%) | 190 (38.5%) | | | | |
| | ICU provider | | | | | 160 (16.2%) | 83 (16.8%) | | | | |
| | Anesthesia/ENT | | | | | 178 (18.0%) | 105 (21.3%) | | | | |
| | HCW in COVID unit | | | | | 76 (7.7%) | 29 (5.9%) | | | | |
| | Healthcare worker in congregated care setting | | | | | 11 (1.1%) | 4 (0.8%) | | | | |
| | First responder | | | | | 115 (11.6%) | 65 (13.2%) | | | | |

| | | Vijayaraghavan et al. | | Polo et al. (EPICOS) | | Llanos-Cuentas et al. | | Grau-Pujol et al. | | Syed et al. | |
|--------------------------|--|-----------------------|-------------|-------------------------|-------------|-----------------------|--------------|-------------------|-------------|------------------|-------------|
| | | HCQ | Placebo | HCQ ² | Placebo | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo |
| N (randomization) | | 213 | 203 | 231 | 223 | 36 | 32 | 142 | 127 | 154 | 46 |
| Age | | 32.3 (9.65) | 31.8 (8.63) | 38 (18-65) | 38 (18,65) | 39.14 (1.53) | 39.28 (1.72) | 39.6 (11.2) | 40.3 (12.8) | 30.25 (NA) | 31.9 (9.13) |
| Female | | 100 (46.9%) | 97 (47.8%) | 149 (64.5%) | 143 (64.1%) | 20 (55.6%) | 20 (62.5%) | 104 (73.2%) | 93 (73.2%) | 68 (44.1%) | 23 (50%) |
| BMI (kg/m ²) | | | | | | | | | | | |
| Current smoker | | 8 (3.8%) | 9 (4.4%) | | | | | 21 (14.9%) | 17 (13.8%) | 19 (12.3%) | 7 (15.2%) |
| Race/ Ethnicity | White | | | | | | | | | | |
| | Asian | | | | | | | | | | |
| | African American | | | | | | | | | | |
| | Hispanic | | | | | | | | | | |
| Comorb idities | Asthma | 0 (0%) | 0 (0%) | 20 (8.7%) | 9 (4.0%) | 3 (8.3%) | 4 (12.5%) | 5 (3.5%) | 2 (1.6%) | | |
| | Diabetes | 7 (3.3%) | 3 (1.5%) | 1 (0.4%) | 3 (1.3%) | 1 (2.8%) | 0 (0%) | 0 (0%) | 1 (0.8%) | 4 (2.6%) | 3 (6.5%) |
| | Hypertension | 2 (0.9%) | 3 (1.5%) | 4 (1.7%) | 19 (8.5%) | 3 (8.3%) | 2 (6.3%) | 2 (1.4%) | 3 (2.4%) | 7 (4.5%) | 2 (4.3%) |
| | None | | | | | | | | | | |
| Practice Location | Emergency Department | 26 (12.2%) | 18 (8.9%) | 20 (8.7%) | 21 (9.4%) | | | | | | |
| | Internal Medicine ward | 130 (64%) | 130 (61%) | | | | | | | | |
| | ICU/anesthesia | | | | | | | | | | |
| | Labor and delivery | | | | | | | | | | |
| | Ambulance | | | 0 (0%) | 0 (0%) | | | | | | |
| | Congregate care setting | | | | | | | | | | |
| | ICU | 53 (24.9%) | 53 (26.1%) | 17 (7.4%) | 13 (5.8%) | | | | | | |
| | Operating room | | | | | | | | | | |
| | EMS, Fire and Police First Responders | | | | | | | | | | |
| Occupation | Nurse | 67 (31.5%) | 68 (33.5%) | 67 (29.0%) | 72 (32.3%) | 6 (16.7%) | 5 (15.6%) | 35 (27.8%) | 40 (28.2%) | 20 (13.0%) | 9 (19.6%) |
| | Physician | 34 (16%) | 31 (15.3%) | 74 (32%) | 66 (29.6%) | 23 (63.9%) | 16 (50%) | 67 (47.2%) | 53 (42.1%) | 118 (76.6%) | 25 (54.3%) |
| | Certified Nurse Assistant | | | | | 1 (2.8%) | 0 (0%) | 12 (8.5%) | 12 (9.5%) | | |
| | ED Technician | | | | | | | | | | |
| | Respiratory therapist | | | | | | | | | | |
| | Nurse or Physician | | | | | | | | | | |
| | Emergency Medicine Provider | | | | | | | | | 2 (1.3%) | 0 (0%) |
| | ICU provider | | | | | | | | | | |
| | Anesthesia/ENT | | | | | | | | | | |
| | HCW in COVID unit | | | | | | | | | | |
| | Healthcare worker in congregate care setting | | | | | | | | | | |
| | First responder | | | | | | | | | 2 (1.3%) | 0 (0%) |

HCQ=Hydroxychloroquine ; ITT= Intention-to-treat ; BMI=Body mass index ; ICU=Intensive care unit; ED=Emergency department ; ENT=Ear, nose, throat ; HCW=Healthcare worker

¹ More than one HCQ groups with different doses are lumped.

² The Polo et al. study randomized participants to four treatment groups, and the HCQ and control groups are used in our meta-analysis.

³ Median (range)

⁴ Median (IQR)

⁵ Mean (range)

⁶ Current or previous smoker




eTable 6. Results of outcome measures in trials included in the meta-analysis. Sample size and the number of participants who had each outcome are reported with proportions (%) in parentheses.

| | Treatment | N (ITT) | Confirmed COVID-19 | Suspected with COVID compatible symptoms | Adverse event ² |
|------------------------------------|------------------|---------|--------------------|--|----------------------------|
| Naggie et al. (HERO-HCQ) | HCQ | 683 | 3 (0.4) | 38 (5.6) | 16 (2.3) |
| | Placebo | 676 | 6 (0.9) | 47 (7.0) | 13 (1.9) |
| Abella et al. (PATCH) | HCQ | 64 | 4 (6.3) | | 29 (45.3) |
| | Placebo | 61 | 4 (6.6) | | 17 (27.9) |
| Rajasingham et al. (MN-COVID-PREP) | HCQ ¹ | 989 | 11 (1.1) | 47 (4.8) | 316 (32.0) |
| | Placebo | 494 | 6 (1.2) | 33 (6.7) | 100 (20.2) |
| Rojas-Serrano et al. | HCQ | 62 | 1 (1.6) | | 32 (51.6) |
| | Placebo | 65 | 6 (9.2) | | 38 (58.5) |
| McKinnon et al. (WHIP) | HCQ ¹ | 387 | 2 (0.5) | 22 (5.7) | 192 (49.6) |
| | Placebo | 191 | 2 (1.0) | 13 (6.8) | 85 (44.5) |
| Vijayaraghavan et al. | HCQ | 211 | 11 (5.2) | 12 (5.7) | 21 (10.0) |
| | Placebo | 203 | 12 (5.9) | 12 (5.9) | 14 (6.9) |
| Polo et al. (EPICOS) | HCQ | 224 | 21 (9.4) | | 100 (44.6) |
| | Placebo | 211 | 23 (10.9) | | 94 (44.5) |
| Llanos-Cuentas et al. | HCQ | 34 | 5 (14.7) | | |
| | Placebo | 31 | 3 (9.7) | | |
| Grau-Pujol et al. | HCQ | 137 | 1 (0.7) | 3 (2.2) | 53 (38.7) |
| | Placebo | 116 | 1 (0.9) | 3 (2.6) | 42 (36.2) |
| Syed et al. | HCQ ¹ | 154 | 42 (27.3) | | 9 (5.8) |
| | Placebo | 46 | 7 (15.2) | | 1 (2.2) |

HCQ= Hydroxychloroquine; ITT= Intention-to-treat; AE=Adverse event ; COVID-RS=COVID-19 related symptoms ; Vit C= Vitamin C

¹ More than one HCQ groups with different doses are lumped.

² Number of patients with any adverse events

eFigure. Funnel plots for the three outcomes**(a) Lab-confirmed positive COVID-19****(b) Suspected COVID-19****(c) Adverse events**



PRISMA 2020 Checklist

| Section and Topic | Item # | Checklist item | Location where item is reported |
|-------------------------------|--------|--|---------------------------------|
| TITLE | | | |
| Title | 1 | Identify the report as a systematic review. | 1 |
| ABSTRACT | | | |
| Abstract | 2 | See the PRISMA 2020 for Abstracts checklist. | 4 |
| INTRODUCTION | | | |
| Rationale | 3 | Describe the rationale for the review in the context of existing knowledge. | 5-6 |
| Objectives | 4 | Provide an explicit statement of the objective(s) or question(s) the review addresses. | 6 |
| METHODS | | | |
| Eligibility criteria | 5 | Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses. | 7 |
| Information sources | 6 | Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted. | 7 |
| Search strategy | 7 | Present the full search strategies for all databases, registers and websites, including any filters and limits used. | 7 |
| Selection process | 8 | Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process. | 7 |
| Data collection process | 9 | Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process. | 7 |
| Data items | 10a | List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect. | 8 |
| | 10b | List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information. | 8 |
| Study risk of bias assessment | 11 | Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process. | 8 |
| Effect measures | 12 | Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results. | 9 |
| Synthesis methods | 13a | Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)). | Supplement |
| | 13b | Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions. | 9 |
| | 13c | Describe any methods used to tabulate or visually display results of individual studies and syntheses. | 8-9 |
| | 13d | Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used. | 10 |
| | 13e | Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression). | 10 |
| | 13f | Describe any sensitivity analyses conducted to assess robustness of the synthesized results. | 10 |
| Reporting bias assessment | 14 | Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases). | 9 |
| Certainty assessment | 15 | Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome. | 9 |



PRISMA 2020 Checklist

| Section and Topic | Item # | Checklist item | Location where item is reported |
|--|--------|--|---------------------------------|
| RESULTS | | | |
| Study selection | 16a | Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram. | 11 |
| | 16b | Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded. | 11-12 |
| Study characteristics | 17 | Cite each included study and present its characteristics. | 8-9, Supplement |
| Risk of bias in studies | 18 | Present assessments of risk of bias for each included study. | Supplement |
| Results of individual studies | 19 | For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots. | Supplement |
| Results of syntheses | 20a | For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies. | Supplement |
| | 20b | Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect. | 11-13 |
| | 20c | Present results of all investigations of possible causes of heterogeneity among study results. | 11-13 |
| | 20d | Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results. | 11-13 |
| Reporting biases | 21 | Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed. | Supplement |
| Certainty of evidence | 22 | Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed. | Supplement |
| DISCUSSION | | | |
| Discussion | 23a | Provide a general interpretation of the results in the context of other evidence. | 14 |
| | 23b | Discuss any limitations of the evidence included in the review. | 16 |
| | 23c | Discuss any limitations of the review processes used. | 16 |
| | 23d | Discuss implications of the results for practice, policy, and future research. | 16 |
| OTHER INFORMATION | | | |
| Registration and protocol | 24a | Provide registration information for the review, including register name and registration number, or state that the review was not registered. | Supplement |
| | 24b | Indicate where the review protocol can be accessed, or state that a protocol was not prepared. | 7 |
| | 24c | Describe and explain any amendments to information provided at registration or in the protocol. | 7 |
| Support | 25 | Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review. | 17 |
| Competing interests | 26 | Declare any competing interests of review authors. | 17 |
| Availability of data, code and other materials | 27 | Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review. | Supplement |

BMJ Open

Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in healthcare workers: a meta-analysis of randomized clinical trials

| | |
|---------------------------------|--|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2022-065305.R2 |
| Article Type: | Original research |
| Date Submitted by the Author: | 09-May-2023 |
| Complete List of Authors: | Hong, Hwanhee; Duke University, Department of Biostatistics and Bioinformatics Friedland, Anne; UNC School of Medicine Hu, Mengyi; Duke University, Department of Biostatistics and Bioinformatics Anstrom, Kevin J.; University of North Carolina System, Collaborative Studies Coordinating Center Halabi, Susan; Duke University School of Medicine, Department of Biostatistics & Bioinformatics McKinnon, John; Henry Ford Hospital, Division of Infectious Diseases Amaravadi, Ravi; University of Pennsylvania Rojas-Serrano, Jorge; Instituto Nacional de Enfermedades Respiratorias, Interstitial Lung Disease and Rheumatology Units Abella, Benjamin; University of Pennsylvania Portillo-Vázquez, Angélica Margarita; Instituto Nacional de Enfermedades Respiratorias Woods, Christopher; Duke Clinical Research Institute Hernandez, Adrian F.; Duke University School of Medicine Boulware, David R.; University of Minnesota Twin Cities Naggie, Susanna; Duke Clinical Research Institute, Infectious Disease Rajasingham, Radha; University of Minnesota Twin Cities |
| Primary Subject Heading: | Infectious diseases |
| Secondary Subject Heading: | Infectious diseases, Evidence based practice |
| Keywords: | COVID-19, STATISTICS & RESEARCH METHODS, EPIDEMIOLOGY |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in
healthcare workers: a meta-analysis of randomized clinical trials**

Hwanhee Hong, PhD (0000-0002-3736-6327), Anne Friedland, MD (0000-0002-9735-4838),
Mengyi Hu, MB (0000-0003-1999-6672), Kevin J. Anstrom, PhD (0000-0001-6452-2172),
Susan Halabi, PhD (0000-0003-4135-2777), John E. McKinnon, M.D., M.Sc (0000-0003-3916-
4021), Ravi Amaravadi, MD (0000-0002-5768-2474), Jorge Rojas Serrano, MD, PhD (0000-
0001-6980-7898), Benjamin S. Abella, MD MPhil (0000-0003-2521-0891), Angélica Margarita
Portillo-Vázquez, MD, MSc (0000-0002-5032-4076), Christopher W. Woods, MD, MPH,
Adrian Hernandez, MD, MHS (0000-0003-3387-9616), David R Boulware, MD MPH (0000-
0002-4715-0060), Susanna Naggie, MD, MHS (0000-0001-7721-6975), Radha Rajasingham,
MD (0000-0001-5531-0231)

Hwanhee Hong, Assistant Professor, Department of Biostatistics and Bioinformatics, Duke
University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA

Anne Friedland, Assistant Professor, Department of Infectious Disease, University of North
Carolina School of Medicine, 130 Mason Farm Rd, Chapel Hill, NC, 27514, USA

Mengyi Hu, Statistician, Department of Biostatistics and Bioinformatics, Duke University, 2424
Erwin Road, Ste 1105, Durham, NC, 27705, USA

Kevin J. Anstrom, Director, Collaborative Studies Coordinating Center, Gillings School of
Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA

Susan Halabi, Professor, Department of Biostatistics and Bioinformatics, Duke University, 2424
Erwin Road, Suite 11088, Durham, NC, 27705, USA

1
2
3 23 John E. McKinnon, Senior Staff, Infectious Diseases, Henry Ford Health, Clara Ford Pavilion,
4
5 24 Suite 322, 2799 W. Grand Blvd., Detroit, MI 48202, USA
6
7 25 Ravi Amaravadi, Professor of Medicine, University of Pennsylvania , 852 BRB 2/3 421 Curie
8
9 26 Blvd., Philadelphia, PA 19063, USA
10
11
12 27 Jorge Rojas Serrano, Attending physician, Interstitial Lung Disease and Rheumatology Unit,
13
14 28 Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas, Calzada de Tlalpan
15
16 29 4502, Tlalpan, Ciudad de México, México. 14080
17
18
19 30 Benjamin S. Abella, Professor, Department of Emergency Medicine, University of Pennsylvania,
20
21 31 423 Guardian Drive Room 412, Philadelphia, PA 19104, USA
22
23 32 Angélica Margarita Portillo-Vázquez, Attending physician, Otolaryngology department, Instituto
24
25 33 Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas. Calzada de Tlalpan 4502,
26
27 34 Tlalpan, Ciudad de México, México. 14080
28
29
30 35 Christopher W. Woods, Professor of Medicine, Duke University, 310 Trent Drive, Durham, NC
31
32 36 27710, USA
33
34
35 37 Adrian Hernandez, Professor of Medicine, Duke Clinical Research Institute, Duke University
36
37 38 300 W. Morgan Street Durham, NC 27701, USA
38
39
40 39 David R Boulware, Professor, Division of Infectious Diseases & International Medicine,
41
42 40 Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis, MN, 55455,
43
44 41 USA
45
46 42 Susanna Naggie, Professor of Medicine, Duke Clinical Research Institute, 300 West Morgan
47
48 43 Street, 5th Floor, Durham, NC 27710, USA
49
50
51
52
53
54
55
56
57
58
59
60

44 Radha Rajasingham, Assistant Professor, Division of Infectious Diseases & International
45 Medicine, Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis,
46 MN, 55455, USA

47
48 **Corresponding author:** Hwanhee Hong, Assistant Professor, Department of Biostatistics and
49 Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA,
50 hwanhee.hong@duke.edu

51
52 Abstract word count: 226

53 Word count: 2475

54
55 **Key words:** SARS-CoV-2, COVID-19, hydroxychloroquine, prophylaxis, health care workers,
56 meta-analysis, clinical trials

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective: We studied the safety and efficacy of hydroxychloroquine (HCQ) as pre-exposure prophylaxis for COVID-19 in healthcare workers (HCWs), using a meta-analysis of randomized controlled trials.

Data Sources: PubMed, and EMBASE databases were searched to identify randomized trials studying HCQ.

Study Selection: Ten randomized controlled trials (RCTs) were identified (n=5,079 participants).

Data Extraction and Synthesis: The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis between HCQ and placebo using a Bayesian random-effects model. A *pre-hoc* statistical analysis plan was written, and the review protocol was registered at PROSPERO (CRD42021285093)

Main Outcomes: The primary efficacy outcome was polymerase chain reaction (PCR)-confirmed SARS-CoV-2 infection and the primary safety outcome was incidence of adverse events. The secondary outcome included clinically suspected SARS-CoV-2 infection.

Results: Compared with placebo, HCWs randomized to hydroxychloroquine (HCQ) had no significant difference in PCR-confirmed SARS-CoV-2 infection (odds ratio [OR] 0.92, 95% credible interval [CI]: 0.58, 1.37) or clinically suspected SARS-CoV-2 infection (OR 0.78, 95% CI: 0.57, 1.10), and marginally significant difference in adverse events (OR 1.35, 95% CI: 1.03, 1.73).

Conclusions and Relevance: Our meta-analysis of ten RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in HCWs found that compared with placebo HCQ

does not significantly reduce the risk of confirmed or clinically suspected SARS-CoV-2 infection, while HCQ significantly increases adverse events.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- Bayesian meta-analysis models with random effects fitted the data.
- The ten trials included in the meta-analysis represent wide geographical locations including US, Canada, Mexico, India, Spain, Bolivia, Venezuela, Peru, and Pakistan.
- The findings can be applied to healthcare workers but should not be generalized to a broader population.

INTRODUCTION

Early during the SARS-CoV-2 pandemic, based on *in vitro* antiviral activity of both chloroquine and hydroxychloroquine against SARS-CoV-2 [1-3], clinicians considered use of hydroxychloroquine (HCQ) for treatment and prevention of SARS-CoV-2 infection and the associated disease, COVID-19. While there are now published randomized controlled trials of HCQ for the treatment of COVID-19 in the inpatient and outpatient setting [4, 5], there remains a lack of adequately powered randomized controlled trials of HCQ for the pre-exposure prophylaxis (PrEP) of SARS-CoV-2 infection. A number of COVID-19 clinical studies including PrEP studies were planned early in the pandemic; however, several never opened to enrollment and those that did open were closed early without reaching full accrual due to the rapidly changing landscape of preventative therapies, including vaccines, and a significant shift in public opinion of HCQ as a medical intervention for SARS-CoV-2 [6].

1
2
3 111 Vaccination access remains insufficient globally [7]. Specifically, in low-income
4
5 112 countries only 33% of healthcare workers are fully vaccinated. While high-income countries
6
7 113 have better coverage, overall 38% of countries did not achieve the milestone of 70% vaccination
8
9
10 114 coverage for healthcare workers by the end of 2021[8]. Thus, studying the pre-exposure
11
12 115 prophylaxis potential for a drug with a known safety profile is crucial to protect people at high
13
14 116 risk of exposures, such as healthcare workers (HCWs) [9, 10]. Two large randomized, placebo-
15
16
17 117 controlled trials testing the safety and efficacy of HCQ as pre-exposure prophylaxis for COVID-
18
19 118 19 in HCWs [11] [12], showed potential for a modest benefit of HCQ but were both
20
21 119 underpowered, if a modest effect exists. More trials [13-15] studying HCQ as pre-exposure
22
23 120 prophylaxis of COVID-19 in HCWs have been published with similar limitations.

24
25
26 121 To address the most common limitation, inadequate power to show a modest effect, we
27
28 122 conducted a formal meta-analysis of pre-exposure prophylactic HCQ studies in HCWs. We
29
30 123 conducted a systematic search for clinical trials of pre-exposure prophylactic use of HCQ against
31
32 124 infection of SARS-CoV-2 in HCWs, thoroughly compared similarities and differences in
33
34 125 characteristics of the identified studies and performed a Bayesian meta-analysis to combine
35
36 126 results of the trials.

37
38 127
39
40
41
42 128 **METHODS**

43
44 129 The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines
45
46 130 were used in this systematic review and meta-analysis[16]. A statistical analysis plan was written
47
48 131 in advance and the review protocol was registered at PROSPERO (CRD42021285093).

49
50 132
51
52
53
54 133 **Search strategy and information sources**

We searched PubMed/Medline and Ovid/Embase databases from database inception through the final search date March 14, 2023. We used keywords related to COVID-19, HCQ, and randomized controlled trials. The full search strategies are provided in eTable 1.

Eligibility criteria and study selection

The eligibility criteria included phase II or phase III randomized controlled trials (RCTs) of hydroxychloroquine for use as pre-exposure prophylaxis in HCWs with moderate to high risk of exposure. We excluded observational studies, crossover trials, studies where the method of allocation to treatment was not truly random, duplicate studies, and non-original data studies. No language, publication date, or publication status restrictions were applied. References of prior systematic reviews and meta-analyses were also screened for related studies. Study selection involved screening of titles and abstracts followed by full-text evaluation of possible eligible studies.

Data collection process

Each of the selected studies were independently reviewed by two reviewers (AF, MH, or HH). We extracted data on the study design, baseline characteristics, interventions, and outcomes. Any disagreements of collected information between reviews were reconciled through discussion by all three reviewers.

Outcome measures

The primary efficacy outcome for the meta-analysis was laboratory confirmed SARS-CoV-2 infection by polymerase chain reaction (PCR) test and the primary safety outcome was incidence

of adverse events (Table 1). The secondary efficacy outcome was suspected or probable SARS-CoV-2 infection. Included studies had the following outcome definitions: (1) laboratory confirmed SARS-CoV-2 infection defined as COVID-19 like symptoms and positive SARS-CoV-2 PCR and (2) suspected or probable SARS-CoV-2 infection defined as COVID-19 like symptoms but lack of confirmatory PCR testing.

Table 1. Treatment strategies, adherence, trial-defined primary outcome, and study duration for trials included in the meta-analysis

| | Trial-defined primary outcome | Study duration | Treatment group | Randomized treatment assignment | Randomized sample size |
|--|--|---------------------|------------------|--|------------------------|
| Naggie et al.[13] (HERO-HCQ) | Confirmed (by NP swab PCR) or suspected COVID-19 infection through 30 days | 60 days | HCQ | HCQ 600 mg BID loading dose for Day 1, followed by 400 mg QD for 29 days | 683 |
| | | | Control | Placebo | 676 |
| Abella et al.[11] (PATCH) | COVID-19 infection as determined by positive NP swab over 8 weeks | 56 days (8 weeks) | HCQ | HCQ 600mg daily for 60 days | 64 |
| | | | Control | Placebo | 61 |
| Rajasingham et al.[12] (MN-COVID-PREP) | COVID-19 free survival time by lab confirmed or probable illness | 84 days (12 weeks) | HCQ ^a | HCQ loading doses (400 mg twice 6-8hrs apart), followed by 400 mg once weekly or 400 mg twice weekly for 84 days | 989 |
| | | | Control | Placebo | 494 |
| Rojas-Serrano et al.[14] | Time to symptomatic respiratory infection with a positive COVID RT PCR over 60 days | 60 days | HCQ | HCQ 200 mg daily for 60 days | 62 |
| | | | Control | Placebo | 65 |
| McKinnon et al.[15] (WHIP) | Lab confirmed cases of COVID-19 determined by either IgM and IgG serology in blood sample or RT-PCR test results | 56 days (8 weeks) | HCQ ^a | HCQ 400 mg loading dose for Day 1, followed by 200 mg daily or 400 mg weekly on the same day of each week for 56 days | 387 |
| | Confirmed new cases of COVID-19 | | Control | Placebo | 191 |
| Vijayaraghavan et al.[17] | Lab confirmed SARS-CoV-2 infection by PCR or presence of antibodies | 180 days (6 months) | HCQ | HCQ 400 mg twice on the day of enrollment, followed by 400 mg once a week for a total of 12 weeks plus personal protective equipment (PPE) | 213 |
| | | | Control | PPE | 203 |
| Polo et al.[18] (EPICOS) | Lab confirmed symptomatic COVID-19 by PCR | 84 days (12 weeks) | HCQ ^b | HCQ 200 mg once daily | 231 |
| | | | Control | Placebo | 223 |

| | | | | | |
|---------------------------|--|---------------------|------------------|---|-----|
| Llanos-Cuentas et al.[19] | COVID-19 cases confirmed by PCR or serological test | 28 days (4 weeks) | HCQ | HCQ loading dose of 600 mg on the first day, followed by 400 mg every other day plus PPE | 36 |
| | | | Control | PPE | 32 |
| Grau-Pujol et al.[20] | COVID-19 confirmed cases with seroconversion or PCR test | 180 days (6 months) | HCQ | HCQ 400 mg daily for the four consecutive days, followed by 400 mg weekly | 142 |
| | | | Control | Placebo | 127 |
| Syed et al.[17] | COVID-19-free survival (COVID-19 confirmed by PCR) | 84 days (12 weeks) | HCQ ^a | HCQ 400 mg twice for Day 1, followed by 400 mg weekly or HCQ 400 mg once every 3 weeks or HCQ 200 mg once every 3 weeks | 154 |
| | | | Control | Placebo | 46 |

HCQ=Hydroxychloroquine

^a More than one HCQ groups with different doses are lumped.

^b The Polo et al. study randomized participants to four treatment groups, and the HCQ and control groups are used in our meta-analysis.

Treatment assignment

Our meta-analysis did not study HCQ dosing specific effects. For studies randomizing participants to more than one HCQ arm with different doses, all HCQ arms were merged and considered as a single HCQ arm. Such studies include the Rajasingham et al., McKinnon et al. and Syed et al. studies.

Risk of bias and certainty of evidence assessment

Two independent reviewers (AF, HH) assessed the risk of bias (low, intermediate, high) of the included studies using the Cochrane's Collaboration tool [21] (eTable 2). We assessed the certainty of evidence using the Grades of Recommendation Assessment, Development and Evaluation (GRADE) approach [22].

Statistical analysis

1
2
3 184 Bayesian logistic regression meta-analysis models under two assumptions (fixed effect and
4
5 185 random effects) were fitted to estimate the odds ratio of having an outcome between
6
7 186 hydroxychloroquine and placebo [23]. The fixed effect model assumes that the odds ratio is
8
9
10 187 constant across studies, while the random effects model accounts for heterogeneity in the odds
11
12 188 ratios across studies. To assess and compare the goodness-of-fit of the fitted fixed and random
13
14 189 effects models, we calculated the Watanabe-Akaike information criterion [24]. In the Bayesian
15
16 190 models, we assigned non-informative prior distributions as no prior information was available.
17
18 191 The odds ratios and the associated 95% credible intervals were estimated using Markov chain
19
20 192 Monte Carlo (MCMC) algorithms. In addition, we calculated Bayesian posterior probabilities of
21
22 193 the odds ratio smaller than 1 or 0.5 for the primary efficacy outcome, and greater than 2 for the
23
24 194 safety outcome [25]. The standard deviation of the random effects and I^2 [26] were estimated to
25
26 195 quantify the between-study heterogeneity, where small values of both metrics indicate slight
27
28 196 heterogeneity. To identify publication bias, we plotted and assessed funnel plots for their
29
30 197 symmetry, and conducted the Egger's test[27]. All Bayesian meta-analyses were conducted
31
32 198 using the `rstan` package (version 2.21.2)[28] in R 4.0.2 [29]. We used two parallel chains,
33
34 199 where each chain consists of 50,000 samples after a 25,000-sample burn-in. We checked
35
36 200 convergence of the MCMC chains for all model parameters using trace plots and Gelman-Rubin
37
38 201 diagnostic statistics [30].
39
40
41
42
43
44
45
46

47 203 **Patient and public involvement**

48
49 204 No patient involved.
50
51
52
53

54 206 **RESULTS**
55
56
57
58
59
60

207 Search results

208 Our database search resulted in 350 unique studies after excluding duplicates. Of those, 339
209 studies were screened out due to irrelevance based on title and abstract screening. Eleven studies
210 were assessed in full-text for eligibility (Figure 1). Of those, one trial was excluded from the
211 meta-analysis because it studied with non-healthcare worker populations. As a result, a total of
212 ten studies in a population consisting of HCWs were identified (Table 1).

214 Study and patient characteristics

215 Study design, population, treatment strategies, and key characteristics are presented in Table 1
216 and eTable 3. A total of 5,079 randomized participants (2,961 randomized to HCQ) from the 10
217 studies were included in the meta-analysis. The ten studies defined HCWs broadly and included
218 first responders (emergency medical services, fire, and police). The follow-up duration of the 10
219 studies ranged from 28 days to 180 days. The HCQ dosing scheme varied across studies,
220 including daily dosing ranging from 200 to 600mg daily with or without a loading dose and once
221 or twice weekly or once every three weeks dosing. The duration of therapy also varied across
222 studies (Table 1). The trial-specific definitions of primary outcome and adverse events are
223 comparable across trials (Table 1, eTable 4).

225 Baseline characteristics by randomized treatment assignment are reported (eTable 5). The
226 average age ranged between 31 and 45. The aggregate proportion of women within each study
227 varied across the 10 trials, with a range from 44% to 69%. In addition, the Abella et al. and
228 Rojas-Serrano et al. studies had smaller sample size compared with the other three studies and
229 showed a difference in female ratio between placebo and HCQ groups. In the Naggie et al.,

1
2
3 230 Abella et al., Rajasingham et al., and McKinnon et al., studies, over 80% of study participants
4
5 231 were white. The Abella et al. and Rajasingham et al. studies had high proportions of HCWs
6
7 232 working in an emergency department (56% and 41%, respectively) and the Abella et al. study
8
9 233 had a high proportion of nurses (67%).
10
11
12 234
13
14 235 Several studies reported treatment adherence assessed by two methods: self-reported adherence
15
16 236 and/or pill count at the end of the study. The Rajasingham et al. study additionally conducted
17
18 237 remote blood sampling to verify HCQ concentrations in a subset. Adherence varied significantly
19
20 238 across the studies, with a low proportion of approximately 52% in the Rojas-Serrano et al. study
21
22 239 and 97-98% in the Abella et al. study.
23
24
25
26 240

27
28 241 **Results of meta-analysis**

29
30 242 Overall, 3.4% (171/5039) developed PCR-confirmed SARS-CoV-2 infection and 5.6% (230/4087)
31
32 243 developed suspected COVID-19 that was not laboratory confirmed. Since the goodness-of-fit
33
34 244 assessment using Watanabe-Akaike information criterion concluded that the random effects meta-
35
36 245 analysis model was as good as or better than the fixed effect meta-analysis model for all outcomes,
37
38 246 we reported the results under the random effects model. Compared with placebo, HCWs
39
40 247 randomized to HCQ had numerically lower rate of PCR-confirmed SARS-CoV-2 infection cases
41
42 248 (odds ratio [OR] 0.92, 95% credible interval [CI]: 0.58, 1.37; GRADE score: moderate certainty),
43
44 249 and suspected or probable SARS-CoV-2 infection cases (OR 0.78, 95% CI: 0.57, 1.10; GRADE
45
46 250 score: moderate certainty). None of these odds ratios were statistically significant. Participants
47
48 251 treated with HCQ had a numerically higher rate of adverse events (OR 1.35, 95% CI: 1.03, 1.73;
49
50 252 GRADE score: moderate certainty) with marginally statistical significance (Figure 2). The
51
52
53
54
55
56
57
58
59
60

outcome data used in our analyses are presented in eTable 6. The summary of GRADE score assessment is provided in eTable 7.

The Bayesian posterior probabilities of the odds ratio less than 1 for the confirmed SARS-CoV-2 infection outcome (i.e., the probability of HCQ favoring over placebo) was 0.67, while the posterior probability of odds ratio less than 0.5 (i.e., the probability that the odds of having a confirmed SARS-CoV-2 infection outcome in HCQ is less than a half of the odds in placebo) was 0.009. The posterior probability of the odds ratio greater than 2 for the adverse event outcome (i.e., the probability that the odds of having an adverse event in HCQ is greater than twice of the odds in placebo) was 0.004.

Our meta-analysis showed little or moderate variability of effect estimates across studies with I^2 value of 0%, 0%, and 43%, and the estimated standard deviation of the random effects of 0.39, 0.26, and 0.45 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively. Funnel plots (eFigure) showed no indication of publication bias and the associated Egger's test results supported that the funnel plots were not asymmetry with p-values of 0.308, 0.305, and 0.794 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively.

DISCUSSION

Understanding the pre-exposure prophylactic effect of HCQ against COVID-19 remains relevant, as its use continues, particularly in the international setting [31, 32]. Our meta-analysis of the ten RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in

1
2
3 276 5,079 HCWs found that HCQ did not have a statistical association with fewer confirmed or
4
5 277 suspected/probable SARS-CoV-2 infection cases compared with placebo. The geographical
6
7 278 locations of the 10 trials included in the meta-analysis are US, Canada, Mexico, India, Spain,
8
9
10 279 Bolivia, Venezuela, Peru, and Pakistan (eTable 3). While the odds ratios of most studies favor
11
12 280 HCQ, the credible intervals remain wide suggesting low certainty in the true point estimate. Two
13
14 281 studies including the Llanos-Cuentas et al. study conducted in Peru and the Syed et al. study
15
16 282 conducted in Pakistan showed odds ratios favoring placebo, though the credible intervals remain
17
18 283 wide. Furthermore, in this population, COVID-19 events rates were low, particularly for the
19
20 284 most relevant PCR-confirmed infection outcome. The low event rate raises further concern for
21
22 285 the uncertainty of these outcomes. Thus, if there is a minimal effect, the absolute benefit would
23
24 286 be low. To gain more certainty, a very large study would need to be done and this is difficult to
25
26 287 support now due to availability of highly effective vaccines. The safety profile of HCQ in the
27
28 288 outpatient setting is well understood [33]. In these outpatient studies there was marginally
29
30 289 statistically significant difference in adverse events in the HCQ versus the placebo arm,
31
32 290 indicating that HCQ is less safe than placebo.
33
34 291
35
36
37
38
39 292 Our findings can be applied to HCWs but should not be generalized to a broader population. Our
40
41 293 systematic search found only one published RCT of pre-exposure prophylaxis for non-healthcare
42
43 294 worker populations and the study were excluded from our meta-analysis. This study was
44
45 295 conducted in Singapore [34] and showed a significant reduction in the risk of COVID-19
46
47 296 infection in the HCQ arm when compared with the comparator arm, vitamin C. However, this
48
49 297 study showed moderate risk of bias as it used an open-label cluster-randomization design, the
50
51 298 Institutional Review Board excluded higher risk persons from the hydroxychloroquine arm only,
52
53
54
55
56
57
58
59
60

and the participants may not be representative of a general population due to the communal living environment.

A Bayesian meta-analysis approach was used to fit the data. The Bayesian meta-analysis approach has several advantages. First, its flexibility and the MCMC sampling methods to estimate posterior distributions provide probability-based quantities (e.g., posterior probability of an odds ratio smaller than 0.5) that complement typical meta-analysis results (e.g., odds ratios and the associated credible intervals) and help decision making [35]. Second, the Bayesian meta-analysis model with random effects estimates the between-study variability better than the frequentist counterparts [36]. Third, when it comes to with binary outcomes, the Bayesian approach handles rare events better than the frequentist counterparts [23].

A recently published meta-analysis by García-Albéniz et al. [37] investigated pre-exposure (seven RCTs included) and post-exposure (four RCTs included) prophylactic effects of HCQ, but not limited to the HCW population. They found significant pre-exposure prophylactic effects of HCQ on SARS-CoV-2 infection, different from ours. The seven pre-exposure prophylaxis RCTs included in the García-Albéniz et al. meta-analysis consisted of six RCTs that were in our meta-analysis and the aforementioned Singapore study that was excluded from our meta-analysis. Our meta-analysis provides the most up-to-date, systematic, and comprehensive evidence about prophylactic effects of HCQ focusing on the HCW population.

Although a meta-analysis allows for combining evidence from multiple studies in a principled way, our meta-analysis has limitations. First, our analysis did not evaluate effects of different

1
2
3 322 HCQ doses and combined multiple HCQ arms using different doses in three studies. The RCTs
4
5 323 included in our meta-analysis studied varying dosing schemes and a meta-analysis using
6
7 324 aggregate-level data is not a sufficient source to study dosing effects. Second, detailed subgroup
8
9 325 analyses were not conducted due to limited information. Individual-level data are required to
10
11 326 study both dosing and subgroup effects.
12
13
14

15 327
16
17 328 Our meta-analysis of ten RCTs investigating safety and efficacy of HCQ as pre-exposure
18
19 329 prophylaxis in HCWs provides the most up-to-date evidence on HCQ. Although most individual
20
21 330 trials were underpowered and showed null data, integrating the results systematically via meta-
22
23 331 analysis contributes to the scientific literature and provides certain answers to the question. We
24
25 332 found that HCQ does not reduce the risk of confirmed or probable SARS-CoV-2 infection, but
26
27 333 increase risk of adverse events compared with placebo. Hydroxychloroquine should not be used
28
29 334 for pre-exposure prophylaxis in the HCW population.
30
31
32

33 335
34
35 336 **Contributors**
36

37 337 All authors fulfill the ICMJE criteria for authorship. HH, SN, RR, and KJA designed the study.
38
39 338 HH, AF, and MH collected and analyzed the data. HH, SN, and RR wrote the manuscript. SH
40
41 339 and KJA provided statistical review and AF, JEM, RA, JRS, BSA, AMPV, CWW, AH and DRB
42
43 340 provided clinical review. All authors approved and decided to submit the paper for publication.
44
45
46 341 Hwanhee Hong – HH
47
48 342 Anne Friedland – AF
49
50 343 Mengyi Hu – MH
51
52 344 Kevin J. Anstrom – KJA
53
54
55
56
57
58
59
60

345 Susan Halabi – SH

346 John E. McKinnon – JEM

347 Ravi Amaravadi – RA

348 Jorge Rojas Serrano – JRS

349 Benjamin S. Abella – BSA

350 Angélica Margarita Portillo-Vázquez – AMPV

351 Christopher W. Woods – CWW

352 Adrian Hernandez – AH

353 David R Boulware – DRB

354 Susanna Naggie – SN

355 Radha Rajasingham – RR

356 **Funding**

357 This study is funded by the Patient Centered Outcomes Research Institute (PCORI), Contract
358 Number COVID-19-2020-001. The funder had no role in the design, conduct, analysis, or
359 reporting of this study.

360 **Competing interests**

361 All authors except Dr. Abella reported no financial relationship with commercial interest. Dr.
362 Abella have received NIH funds for COVID-19 related research, and holds equity in VOC
363 Health, a start-up company that is developing novel covid testing.

364 **Ethics Approval**

365 Ethics approval was not required because this study used publicly available aggregate data that
366 were not involved with patients' information or prospective data collection.

367 **Data sharing statement**

The data are presented in eTable 6.

REFERENCES

1. Kalil, A.C., *Treating COVID-19—off-label drug use, compassionate use, and randomized clinical trials during pandemics*. JAMA, 2020. **323**(19): p. 1897-1898.

2. McCreary, E.K., J.M. Pogue, and o.b.o.t.S.o.I.D. Pharmacists, *Coronavirus Disease 2019 Treatment: A Review of Early and Emerging Options*. Open Forum Infectious Diseases, 2020. **7**(4).

3. Wang, M., et al., *Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro*. Cell research, 2020. **30**(3): p. 269-271.

4. RECOVERY Collaborative Group, *Effect of hydroxychloroquine in hospitalized patients with Covid-19*. New England Journal of Medicine, 2020. **383**(21): p. 2030-2040.

5. Skipper, C.P., et al., *Hydroxychloroquine in nonhospitalized adults with early COVID-19: a randomized trial*. Annals of internal medicine, 2020. **173**(8): p. 623-631.

6. Halabi, S., et al., *Landscape of coronavirus disease 2019 clinical trials: New frontiers and challenges*. Clinical Trials, 2022: p. 17407745221105106.

7. Padma, T., *COVID vaccines to reach poorest countries in 2023—despite recent pledges*. Nature, 2021. **595**(7867): p. 342-343.

8. Nabaggala, M.S., et al., *The global inequity in COVID-19 vaccination coverage among health and care workers*. International Journal for Equity in Health, 2022. **21**(3): p. 147.

9. World Health Organization. *Prevention, identification and management of health worker infection in the context of COVID-19*. 2020 [cited 2022 May 13th]; Available from: <https://www.who.int/publications/i/item/10665-336265>

10. The United Kingdom Office for National Statistics. *Coronavirus (COVID-19) infections in the community in England: May 2021*. 2021 [cited 2022 May 13th]; Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristicsofpeopletestingpositiveforcovid19incountriesoftheuk20may2021#percentage-testing-positive-for-covid-19-by-patient-facing-and-non-patient-facing-job-roles-uk>.

11. Abella, B.S., et al., *Efficacy and safety of hydroxychloroquine vs placebo for pre-exposure SARS-CoV-2 prophylaxis among health care workers: a randomized clinical trial*. JAMA internal medicine, 2021. **181**(2): p. 195-202.

12. Rajasingham, R., et al., *Hydroxychloroquine as Pre-exposure Prophylaxis for Coronavirus Disease 2019 (COVID-19) in Healthcare Workers: A Randomized Trial*. Clinical Infectious Diseases, 2020. **72**(11): p. e835-e843.

13. Naggie, S., et al., *Hydroxychloroquine for pre-exposure prophylaxis of COVID-19 in health care workers: a randomized, multicenter, placebo-controlled trial Healthcare Worker Exposure Response and Outcomes of Hydroxychloroquine (HERO-HCQ)*. International Journal of Infectious Diseases, 2023. **129**: p. 40-48.

14. Rojas-Serrano, J., et al., *Hydroxychloroquine for prophylaxis of COVID-19 in health workers: A randomized clinical trial*. PLoS One, 2022. **17**(2): p. e0261980.

15. McKinnon, J.E., et al., *Safety and tolerability of hydroxychloroquine in health care workers and first responders for the prevention of COVID-19: WHIP COVID-19 Study*. International Journal of Infectious Diseases, 2022. **116**: p. 167-173.
16. Hutton, B., et al., *The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations*. Annals of internal medicine, 2015. **162**(11): p. 777-784.
17. Tirupakuzhi Vijayaraghavan, B.K., et al., *Hydroxychloroquine plus personal protective equipment versus personal protective equipment alone for the prevention of laboratory-confirmed COVID-19 infections among healthcare workers: a multicentre, parallel-group randomised controlled trial from India*. BMJ Open, 2022. **12**(6): p. e059540.
18. Polo, R., et al., *Daily tenofovir disoproxil fumarate/emtricitabine and hydroxychloroquine for pre-exposure prophylaxis of COVID-19: a double-blind placebo-controlled randomized trial in healthcare workers*. Clinical Microbiology and Infection, 2023. **29**(1): p. 85-93.
19. Llanos-Cuentas, A., et al., *Hydroxychloroquine to prevent SARS-CoV-2 infection among healthcare workers: early termination of a phase 3, randomised, open-label, controlled clinical trial*. BMC Research Notes, 2023. **16**(1): p. 22.
20. Grau-Pujol, B., et al., *Pre-exposure prophylaxis with hydroxychloroquine for COVID-19: a double-blind, placebo-controlled randomized clinical trial*. Trials, 2021. **22**(1): p. 808.
21. Sterne, J.A., et al., *RoB 2: a revised tool for assessing risk of bias in randomised trials*. bmj, 2019. **366**.
22. Puhan, M.A., et al., *A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis*. Bmj, 2014. **349**.
23. Hong, H., C. Wang, and G.L. Rosner, *Meta-analysis of rare adverse events in randomized clinical trials: Bayesian and frequentist methods*. Clinical Trials, 2021. **18**(1): p. 3-16.
24. Watanabe, S. and M. Opper, *Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory*. Journal of machine learning research, 2010. **11**(12).
25. Ferreira, D., et al., *Bayesian predictive probabilities: a good way to monitor clinical trials*. British journal of anaesthesia, 2021. **126**(2): p. 550-555.
26. Higgins, J.P. and S.G. Thompson, *Quantifying heterogeneity in a meta-analysis*. Statistics in medicine, 2002. **21**(11): p. 1539-1558.
27. Egger, M., et al., *Bias in meta-analysis detected by a simple, graphical test*. Bmj, 1997. **315**(7109): p. 629-634.
28. Stan Development Team, *RStan: the R interface to Stan*. R package version, 2020. **2.21.2**.
29. R Core Team, *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2021.
30. Gelman, A. and D.B. Rubin, *Inference from iterative simulation using multiple sequences*. Statistical science, 1992. **7**(4): p. 457-472.
31. Infante, M., et al., *Hydroxychloroquine in the COVID-19 pandemic era: in pursuit of a rational use for prophylaxis of SARS-CoV-2 infection*. Expert review of anti-infective therapy, 2021. **19**(1): p. 5-16.
32. *Revised advisory on the use of hydroxychloroquine (HCQ) as prophylaxis for SARS-CoV-2 infection (in supersession of previous advisory dated 23rd March. 2020)*. 2022; Available from:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[https://www.icmr.gov.in/pdf/covid/techdoc/V5 Revised advisory on the use of HCQ SARS CoV2 infection.pdf](https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ_SARS_CoV2_infection.pdf).

33. Lofgren, S.M., et al. *Safety of hydroxychloroquine among outpatient clinical trial participants for COVID-19*. in *Open forum infectious diseases*. 2020. Oxford University Press US.

34. Seet, R.C.S., et al., *Positive impact of oral hydroxychloroquine and povidone-iodine throat spray for COVID-19 prophylaxis: An open-label randomized trial*. *International Journal of Infectious Diseases*, 2021. **106**: p. 314-322.

35. Hong, H., et al., *A Bayesian missing data framework for generalized multiple outcome mixed treatment comparisons*. *Research synthesis methods*, 2016. **7**(1): p. 6-22.

36. Hong, H., et al., *Comparing Bayesian and frequentist approaches for multiple outcome mixed treatment comparisons*. *Medical Decision Making*, 2013. **33**(5): p. 702-714.

37. García-Albéniz, X., et al., *Systematic review and meta-analysis of randomized trials of hydroxychloroquine for the prevention of COVID-19*. *European Journal of Epidemiology*, 2022. **37**(8): p. 789-796.

Figure Legends

Figure 1. Flowchart of literature review

Figure 2. Forest plots of the meta-analysis results showing the number of events (y), sample size (n), posterior median of odds ratios, and the associated 95% credible intervals comparing HCQ versus placebo for (a) lab-confirmed positive COVID-19, (b) suspected COVID-19, and (c) adverse events.

For peer review only

1 Identification
2
3
4
5

6
7
8
9
10
11


12 Screening
13
14
15
16
17


18
19
20
21
22
23
24
25

26 Eligibility
27
28
29
30
31
32
33


34
35
36
37
38
39

40 Included
41
42
43
44
45
46
47
48






(b) Suspected COVID-19



(c) Adverse events



Supplementary Materials

CONTENTS

- eTable 1. Search code
- eTable 2. Risk of bias
- eTable 3. Characteristics of included trials
- eTable 4. Definition of adverse events
- eTable 5. Baseline characteristics
- eTable 6. Results of outcome measures in each study
- eFigure. Funnel plots for the three outcomes
- eTable 7. Summary of GRADE score findings



















































eTable 1. Search code that was used to identify publications as of March 14, 2023**PubMed search**

| | |
|----|--|
| #1 | covid[Title] OR coronavirus[Title] OR sars-cov-2[Title] |
| #2 | hydroxychloroquine[Title] |
| #3 | randomized[Title/Abstract] OR randomized[Title/Abstract] |
| #4 | #1 AND #2 AND #3 |

Embase search

| | |
|----|---|
| #1 | covid:ti OR coronavirus:ti OR 'sars cov 2':ti |
| #2 | hydroxychloroquine:ti |
| #3 | randomized:ab,ti OR randomised:ab,ti |
| #4 | #1 AND #2 AND #3 |

eTable 2. Risk of bias for trials included in the meta-analysis using the Cochrane risk assessment tool. Green circle is for low risk and yellow circle is for some concerns

| | Selection bias (Randomization process) | Performance bias (Deviations from the intended interventions) | Attrition bias ¹ (Missing outcome data) | Reporting bias (Measurement of the outcome) | Other sources of bias (Selection of the reported result) |
|---------------------------------------|---|---|---|---|---|
| Naggie et al. (HERO-HCQ) |  |  |  |  |  |
| Abella et al. (PATCH) |  |  |  |  |  |
| Rajasingham et al. (MN-COVID-PREP) |  |  |  |  |  |
| Rojas-Serrano et al. |  |  |  |  |  |
| McKinnon et al. (WHIP) |  |  |  |  |  |
| Vijayaraghavan et al. |  |  |  |  |  |
| Polo et al. (EPICOS) |  |  |  |  |  |
| Llanos-Cuentas et al. |  |  |  |  |  |
| Grau-Pujol et al. |  |  |  |  |  |
| Syed et al. |  |  |  |  |  |

¹ The Rojas-Serrano et al. study reported minimal loss to follow-up (<10%). The Rojas-Serrano et al. study reported 18% (25/130) lost to follow-up and additional 12% (16/130) discontinued the intervention.

eTable 3. Characteristics of trials included in the meta-analysis

| | Naggie et al. (HERO-HCQ) | Abella et al. (PATCH) | Rajasingham et al. (MN-COVID-PREP) | Rojas-Serrano et al. | McKinnon et al. (WHIP) |
|---|--|--|--|---|--|
| N (randomization) | 1360 | 132 | 1496 | 130 | 624 |
| Study start date ¹ | 4/22/2020 | 4/9/2020 | 4/6/2020 | 4/21/2020 | 4/10/2020 |
| Study completion date ² | 1/9/2021 | 11/13/2020 | 7/13/2020 | 3/31/2021 | 12/14/2020 |
| Occupation | HCWs at risk of COVID exposure through work in the ICU, emergency department, emergency services, respiratory services or COVID unit | HCWs (Physicians, nurses, certified nursing assistants, emergency technicians, respiratory therapists) eligible working >20 hrs/week | HCWs (physicians, nurses, emergency medical technicians) with direct contact with COVID patients including emergency department and ICU setting, first responders and performing aerosol generating procedures | HCWs (nurses, nursing aids, cleaning staff, orderlies, respiratory therapists and physicians) taking care of hospitalized patients with COVID | HCW, first responders and correlational/law officers, nursing home workers, medical students, public transit workers, household family members of HCW in Michigan and Ohio |
| Sites | 34 sites across the US | 2 tertiary urban hospitals | Multiple sites nationwide across US and Canada | Single site (National Institute of Respiratory Diseases of Mexico) | Multiple sites at Michigan in the US |
| Randomization | Yes (Phase III) | Yes (Phase II) | Yes (Phase III) | Yes (Phase III) | Yes (Phase III) |
| Trial type | Double-blinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | No weight requirement | <40kg excluded | <50kg excluded | N/A |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Excluded | Excluded | Excluded | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Included | Exclude |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Included | |
| Significant renal disease (stage IV or greater) | Excluded | Included | Excluded | Excluded | Exclude |
| Pregnant/breastfeeding | Included | Excluded | Included in US, Excluded in Canada | Excluded | Exclude |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Excluded | Included | Exclude |
| Use of other medications with significant drug interactions | Included | Excluded | Excluded | Included | N/A |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine or azithromycin) | Any treatment for COVID-19 within 14 days excluded | Current use of HCQ or chloroquine excluded | HCQ or chloroquine within 30 days excluded | Chronic use of HCQ included |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | N/A | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Excluded | Excluded if symptoms within 2 weeks unless negative COVID test | Excluded | Excluded | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | N/A |
| Positive COVID serology | Included | Included | N/A | Included | N/A |
| Analysis | Modified intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

| | Vijayaraghavan et al. | Polo et al. (EPICOS) | Llanos-Cuentas et al. | Grau-Pujol et al. | Syed et al. |
|---|---|--|--|--|--|
| N (randomization) | 416 | 454 | 68 | 269 | 200 |
| Study start date ¹ | 6/29/2020 | 4/2020 Spain, 10/2020 Bolivia, 3/2021 Venezuela | June, 2020 | 4/4/2020 | 5/1/2020 |
| Study completion date ² | 2/4/2021 | 5/30/2021 | November, 2020 | Study halted a 1 month analysis | Not reported |
| Occupation | HCWs in an environment with exposure to COVID-19 (physicians, nurses, allied health workers and ancillary health workers) | HCWs (physicians, nurses, medical students, other workers with and without direct patient contact) | HCWs (physicians, nursing staff, technical staff and nursing assistants involved in care of COVID-19 patients) | HCWs (physicians, nurses, nurse assistants and administrators working at least 3 days a week in the trial hospitals) | HCWs at risk of COVID-19 exposure including physicians, nurses, first responders, those performing aerosol generating procedures or working in the emergency department, ICU, and general medicine wards |
| Sites | 9 hospitals across India | Multiple sites across Spain, Venezuela and Bolivia | 4 public hospitals across the Lima metropolitan area | 3 hospitals in Barcelona, Spain | Single hospital in Pakistan |
| Randomization | Yes | Yes | Yes (Phase III) | Yes | Yes (Phase II) |
| Trial type | Unblinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18-70 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | <40kg excluded | No weight requirement | No weight requirement | <40 kg |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Included | Excluded | Included | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Excluded | Excluded |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Excluded | Excluded |
| Significant renal disease (stage IV or greater) | Included | Excluded | Excluded | Excluded | Excluded |
| Pregnant/breastfeeding | Excluded | Excluded | Included | Excluded | Excluded |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Included | Excluded | Excluded |
| Use of other medications with significant drug interactions | Excluded | Included | Included | Excluded | Excluded |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine azithromycin) | Any medication as prophylaxis against COVID-19 after 3/1/21 | Use of hydroxychloroquine, chloroquine or azithromycin in the last 30 days excluded | Treatment with chloroquine or hydroxychloroquine within the last 1 month | Those already taking hydroxychloroquine were excluded |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | Excluded | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Not specified in exclusion criteria | Excluded | Not specified in exclusion criteria | Not specified in exclusion criteria | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | Excluded |
| Positive COVID serology | N/A | N/A | N/A | Excluded | Excluded |
| Analysis | Intention-to-treat | Not reported | Intention-to-treat | Intention-to-treat | Not reported |

HCW=Healthcare workers; ICU=Intensive care unit; ¹ Date when first participant was enrolled; ² Date when final data were collected for the last participant

eTable 4. Definition of adverse events

| Trial | AE definition |
|---|---|
| Naggie et al. (HERO-HCQ) | Adverse events include general disorders and administration site conditions, psychiatric disorders, skin and subcutaneous tissue disorders, cardiac disorders, infections and infestations, nervous system disorders, gastrointestinal disorders, investigations (electrocardiogram QT prolonged and heart rate increased), ear and labyrinth disorders, renal and urinary disorders, and respiratory, thoracic and mediastinal disorders. |
| Abella et al. (PATCH) | Adverse events include abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. |
| Rajasingham et al. (MN-COVID-PREP) | Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and others. |
| Rojas-Serrano et al. | Examples of adverse events are as follows: abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and other. |
| McKinnon et al. (WHIP) | Covid-19 related symptoms, covid-19 clinical disease and medication adverse effects including gastrointestinal disorders, nervous system disorders, respiratory, thoracic and mediastinal disorders, general disorders and administration site conditions, cardiac disorders, musculoskeletal and connective tissue disorders, psychiatric disorders, skin and subcutaneous tissue disorders, ear and labyrinth disorders, and eye disorders. |
| Vijayaraghavan et al. | Adverse events listed in each category at the participant level were categorized as cardiac, gastro-intestinal, headache, and Qtc prolongation. |
| Polo et al. (EPICOS) | Adverse events were classified by organ system and included: gastrointestinal disorders, blood and lymphatic system disorders, cardiac disorders, ear and labyrinth disorders, eye disorder, general disorders, immune system disorder, infections, injuries, investigations, metabolism and nutrition disorders, musculoskeletal/connective tissue disorders, nervous system disorders, psychiatric disorders, renal and urinary disorders, reproductive system disorders, respiratory disorders, skin disorders and vascular disorders. |
| Llanos-Cuentas et al. | Adverse events from grade 1 to grade 3 and above. Note that the Llanos-Cuentas et al. study did report the number of adverse events (not participants) in the HCQ group only. Due to limited information, it was excluded from the meta-analysis with the adverse event outcome. |
| Grau-Pujol et al. | Adverse events included: general symptoms (fever, chills, sweating, malaise, myalgia, arthralgia), gastrointestinal symptoms (nausea, abdominal pain, diarrhea, dysgeusia), dermatological symptoms (itching, rash), respiratory symptoms (rhinorrhea, sore throat / odynophagia, cough, pleuritic pain, dyspnea), neurologic symptoms (headache, visual disturbances), and cardiovascular symptoms. Events were graded mild, moderate and severe. |
| Syed et al. | Syed et al. report the number of patients in each group who experienced adverse events, but did not report what the events were. Due to limited information, it was excluded from the meta-analysis with the adverse event outcome. |

eTable 5. Baseline characteristics with additional variables and detailed information. Sample mean and standard deviation (in parenthesis) are reported for continuous variables, and the number of participants and proportion (in parenthesis) are reported for binary or categorical variables.

| | | Naggie et al. (HERO-HCQ) | | Abella et al. (PATCH) | | Rajasingham et al. (MN-COVID-PREP) | | Rojas-Serrano et al. | | McKinnon et al. (WHIP) | |
|--------------------|--|-----------------------------|--------------------|--------------------------|-------------------------|---------------------------------------|--------------------------|-----------------------------|-------------------------------|--|-------------|
| | | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo |
| N (randomization) | | 683 | 676 | 66 | 66 | 989 | 494 | 62 | 65 | 387 | 191 |
| Age | | 44.2 (11.9) | 43.1 (11.2) | 31 (20-66) ³ | 34 (23-62) ³ | 41.5 (35, 49) ³ | 40 (34, 48) ³ | 31.0 (26.4-39) ⁴ | 31.9 (27.2-43.7) ⁴ | 45.7 (11.6); 44.9 (11.4) ² | 44.1 (12.7) |
| Female | | 442 (64.7%) | 446 (66.0%) | 54 (82%) | 37 (56%) | 519 (52.5%) | 241 (48.8%) | 29 (42.6%) | 42 (64.6%) | 220 (57%) | 114 (60%) |
| BMI (kg/m^2) | | 28.3 (6.3) | 28.6 (6.7) | 26 (19-37) ⁵ | 26 (20-50) ⁵ | | | 26.7 (3.9) | 27.2 (4.6) | | |
| Current smoker | | | | 0 (0%) | 0 (0%) | 38 (3.84%) | 13 (2.6%) | 20 (32.2%) ⁶ | 23 (35.4%) ⁶ | | |
| Race/ Ethnicity | White | 624 (91.4%) | 610 (90.2%) | 55 (83%) | 54 (82%) | 852 (86.1%) | 419 (84.8%) | | | 334 (86%) | 161 (84%) |
| | Asian | | | 7 (11%) | 7 (11%) | 46 (4.7%) | 29 (5.9%) | | | 23 (6%) | 15 (8%) |
| | African American | 18 (2.6%) | 23 (3.4%) | 3 (4%) | 1 (2%) | 10 (1.0%) | 10 (2.0%) | | | 15 (4%) | 9 (5%) |
| | Hispanic | 39 (5.7%) | 40 (5.9%) | 0 (0%) | 2 (3%) | 40 (4.0%) | 18 (3.6%) | | | 11 (3%) | 7 (4%) |
| Comorb idities | Asthma | 58 (8.5%) | 77 (11.4%) | 9 (14%) | 14 (21%) | 91 (9.2%) | 59 (11.9%) | | | | |
| | Diabetes | 20 (2.9%) | 35 (5.2%) | 1 (2%) | 3 (5%) | 36 (3.6%) | 14 (2.8%) | | | | |
| | Hypertension | 99 (14.5%) | 99 (14.6%) | 3 (5%) | 14 (21%) | 145 (14.7%) | 60 (12.1%) | | | | |
| | None | | | 54 (82%) | 40 (61%) | 646 (65.3%) | 336 (68.0%) | 53 (85.5%) | 58 (89.2%) | | |
| Practice Location | Emergency Department | 96 (14.1%) | 94 (13.9%) | 38 (58%) | 36 (55%) | 417 (42.2%) | 190 (38.5%) | | | 48 (12%) | 19 (10%) |
| | Internal Medicine ward | | | 17 (26%) | 18 (27%) | 98 (9.9%) | 56 (11.3%) | | | 31 (8%) | 20 (10%) |
| | ICU/anesthesia | | | 6 (9%) | 6 (9%) | | | | | | |
| | Labor and delivery | | | 5 (7%) | 6 (9%) | | | | | | |
| | Ambulance | 66 (9.7%) | 63 (9.3%) | | | 73 (7.4%) | 45 (9.1%) | | | | |
| | Congregate care setting | | | | | 46 (4.7%) | 20 (4.0%) | | | | |
| | ICU | 48 (7.0%) | 59 (8.7%) | | | 184 (18.6%) | 85 (17.2%) | | | 37 (10%) | 23 (12%) |
| | Operating room | | | | | 103 (10.4%) | 75 (15.2%) | | | | |
| | EMS, Fire and Police | | | | | | | | | 32 (8%) | 16 (8%) |
| | First Responders | | | | | | | | | | |
| Occupation | Nurse | 186/677 (27.5%) | 167/668 (25.0%) | 46 (70%) | 42 (64%) | | | | | | |
| | Physician | 143/677 (21.1%) | 144/668 (21.6%) | 11 (17%) | 16 (24%) | | | | | | |
| | Certified Nurse Assistant | | | 2 (3%) | 2 (3%) | | | | | | |
| | ED Technician | | | 3 (4%) | 1 (2%) | | | | | | |
| | Respiratory therapist | 15/677 (2.2%) | 18/668 (2.7%) | 3 (4%) | 5 (7%) | | | | | | |
| | Nurse or Physician | | | | | | | 31 (50%) | 33 (50.8%) | | |
| | Emergency Medicine Provider | | | | | 407 (41.1%) | 190 (38.5%) | | | | |
| | ICU provider | | | | | 160 (16.2%) | 83 (16.8%) | | | | |
| | Anesthesia/ENT | | | | | 178 (18.0%) | 105 (21.3%) | | | | |
| | HCW in COVID unit | | | | | 76 (7.7%) | 29 (5.9%) | | | | |
| | Healthcare worker in congregare care setting | | | | | 11 (1.1%) | 4 (0.8%) | | | | |
| | First responder | | | | | 115 (11.6%) | 65 (13.2%) | | | | |

| | | Vijayaraghavan et al. | | Polo et al. (EPICOS) | | Llanos-Cuentas et al. | | Grau-Pujol et al. | | Syed et al. | |
|--------------------------|--|-----------------------|-------------|-------------------------|-------------|-----------------------|--------------|-------------------|-------------|------------------|-------------|
| | | HCQ | Placebo | HCQ ² | Placebo | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo |
| | N (randomization) | 213 | 203 | 231 | 223 | 36 | 32 | 142 | 127 | 154 | 46 |
| | Age | 32.3 (9.65) | 31.8 (8.63) | 38 (18-65) | 38 (18,65) | 39.14 (1.53) | 39.28 (1.72) | 39.6 (11.2) | 40.3 (12.8) | 30.25 (NA) | 31.9 (9.13) |
| Female | | 100 (46.9%) | 97 (47.8%) | 149 (64.5%) | 143 (64.1%) | 20 (55.6%) | 20 (62.5%) | 104 (73.2%) | 93 (73.2%) | 68 (44.1%) | 23 (50%) |
| BMI (kg/m ²) | | | | | | | | | | | |
| Current smoker | | 8 (3.8%) | 9 (4.4%) | | | | | 21 (14.9%) | 17 (13.8%) | 19 (12.3%) | 7 (15.2%) |
| Race/ Ethnicity | White | | | | | | | | | | |
| | Asian | | | | | | | | | | |
| | African American | | | | | | | | | | |
| | Hispanic | | | | | | | | | | |
| Comorb idities | Asthma | 0 (0%) | 0 (0%) | 20 (8.7%) | 9 (4.0%) | 3 (8.3%) | 4 (12.5%) | 5 (3.5%) | 2 (1.6%) | | |
| | Diabetes | 7 (3.3%) | 3 (1.5%) | 1 (0.4%) | 3 (1.3%) | 1 (2.8%) | 0 (0%) | 0 (0%) | 1 (0.8%) | 4 (2.6%) | 3 (6.5%) |
| | Hypertension | 2 (0.9%) | 3 (1.5%) | 4 (1.7%) | 19 (8.5%) | 3 (8.3%) | 2 (6.3%) | 2 (1.4%) | 3 (2.4%) | 7 (4.5%) | 2 (4.3%) |
| | None | | | | | | | | | | |
| Practice Location | Emergency Department | 26 (12.2%) | 18 (8.9%) | 20 (8.7%) | 21 (9.4%) | | | | | | |
| | Internal Medicine ward | 130 (64%) | 130 (61%) | | | | | | | | |
| | ICU/anesthesia | | | | | | | | | | |
| | Labor and delivery | | | | | | | | | | |
| | Ambulance | | | 0 (0%) | 0 (0%) | | | | | | |
| | Congregate care setting | | | | | | | | | | |
| | ICU | 53 (24.9%) | 53 (26.1%) | 17 (7.4%) | 13 (5.8%) | | | | | | |
| | Operating room | | | | | | | | | | |
| | EMS, Fire and Police First Responders | | | | | | | | | | |
| Occupation | Nurse | 67 (31.5%) | 68 (33.5%) | 67 (29.0%) | 72 (32.3%) | 6 (16.7%) | 5 (15.6%) | 35 (27.8%) | 40 (28.2%) | 20 (13.0%) | 9 (19.6%) |
| | Physician | 34 (16%) | 31 (15.3%) | 74 (32%) | 66 (29.6%) | 23 (63.9%) | 16 (50%) | 67 (47.2%) | 53 (42.1%) | 118 (76.6%) | 25 (54.3%) |
| | Certified Nurse Assistant | | | | | 1 (2.8%) | 0 (0%) | 12 (8.5%) | 12 (9.5%) | | |
| | ED Technician | | | | | | | | | | |
| | Respiratory therapist | | | | | | | | | | |
| | Nurse or Physician | | | | | | | | | | |
| | Emergency Medicine Provider | | | | | | | | | 2 (1.3%) | 0 (0%) |
| | ICU provider | | | | | | | | | | |
| | Anesthesia/ENT | | | | | | | | | | |
| | HCW in COVID unit | | | | | | | | | | |
| | Healthcare worker in congregate care setting | | | | | | | | | | |
| | First responder | | | | | | | | | 2 (1.3%) | 0 (0%) |

HCQ=Hydroxychloroquine ; ITT= Intention-to-treat ; BMI=Body mass index ; ICU=Intensive care unit; ED=Emergency department ; ENT=Ear, nose, throat ; HCW=Healthcare worker

¹ More than one HCQ groups with different doses are lumped.

² The Polo et al. study randomized participants to four treatment groups, and the HCQ and control groups are used in our meta-analysis.

³ Median (range)

⁴ Median (IQR)

⁵ Mean (range)

⁶ Current or previous smoker




eTable 6. Results of outcome measures in trials included in the meta-analysis. Sample size and the number of participants who had each outcome are reported with proportions (%) in parentheses.

| | Treatment | N (ITT) | Confirmed COVID-19 | Suspected with COVID compatible symptoms | Adverse event ² |
|------------------------------------|------------------|---------|--------------------|--|----------------------------|
| Naggie et al. (HERO-HCQ) | HCQ | 683 | 3 (0.4) | 38 (5.6) | 16 (2.3) |
| | Placebo | 676 | 6 (0.9) | 47 (7.0) | 13 (1.9) |
| Abella et al. (PATCH) | HCQ | 64 | 4 (6.3) | | 29 (45.3) |
| | Placebo | 61 | 4 (6.6) | | 17 (27.9) |
| Rajasingham et al. (MN-COVID-PREP) | HCQ ¹ | 989 | 11 (1.1) | 47 (4.8) | 316 (32.0) |
| | Placebo | 494 | 6 (1.2) | 33 (6.7) | 100 (20.2) |
| Rojas-Serrano et al. | HCQ | 62 | 1 (1.6) | | 32 (51.6) |
| | Placebo | 65 | 6 (9.2) | | 38 (58.5) |
| McKinnon et al. (WHIP) | HCQ ¹ | 387 | 2 (0.5) | 22 (5.7) | 192 (49.6) |
| | Placebo | 191 | 2 (1.0) | 13 (6.8) | 85 (44.5) |
| Vijayaraghavan et al. | HCQ | 211 | 11 (5.2) | 12 (5.7) | 21 (10.0) |
| | Placebo | 203 | 12 (5.9) | 12 (5.9) | 14 (6.9) |
| Polo et al. (EPICOS) | HCQ | 224 | 21 (9.4) | | 100 (44.6) |
| | Placebo | 211 | 23 (10.9) | | 94 (44.5) |
| Llanos-Cuentas et al. | HCQ | 34 | 5 (14.7) | | |
| | Placebo | 31 | 3 (9.7) | | |
| Grau-Pujol et al. | HCQ | 137 | 1 (0.7) | 3 (2.2) | 53 (38.7) |
| | Placebo | 116 | 1 (0.9) | 3 (2.6) | 42 (36.2) |
| Syed et al. | HCQ ¹ | 154 | 42 (27.3) | | 9 (5.8) |
| | Placebo | 46 | 7 (15.2) | | 1 (2.2) |

HCQ= Hydroxychloroquine; ITT= Intention-to-treat; AE=Adverse event ; COVID-RS=COVID-19 related symptoms ; Vit C= Vitamin C

¹ More than one HCQ groups with different doses are lumped.

² Number of patients with any adverse events

eFigure. Funnel plots for the three outcomes**(a) Lab-confirmed positive COVID-19****(b) Suspected COVID-19****(c) Adverse events**

eTable 7. Summary of GRADE score assessment

The summary table is applied to all three outcomes. The GRADE scores for the odds ratios with respect to all three outcomes were downgraded by 1 due to wide credible intervals of odds ratios, resulting in moderate certainty of evidence.

| Item | Quality of evidence |
|------------------|---------------------|
| Risk of bias | High |
| Inconsistency | High |
| Indirectness | High |
| Imprecision | Moderate |
| Publication bias | High |

GRADE Working Group grades of evidence is available here:
<https://gdt.gradeapro.org/app/handbook/handbook.html>



PRISMA 2020 Checklist

| Section and Topic | Item # | Checklist item | Location where item is reported |
|-------------------------------|--------|--|---------------------------------|
| TITLE | | | |
| Title | 1 | Identify the report as a systematic review. | 1 |
| ABSTRACT | | | |
| Abstract | 2 | See the PRISMA 2020 for Abstracts checklist. | 4 |
| INTRODUCTION | | | |
| Rationale | 3 | Describe the rationale for the review in the context of existing knowledge. | 5-6 |
| Objectives | 4 | Provide an explicit statement of the objective(s) or question(s) the review addresses. | 6 |
| METHODS | | | |
| Eligibility criteria | 5 | Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses. | 7 |
| Information sources | 6 | Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted. | 7 |
| Search strategy | 7 | Present the full search strategies for all databases, registers and websites, including any filters and limits used. | 7 |
| Selection process | 8 | Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process. | 7 |
| Data collection process | 9 | Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process. | 7 |
| Data items | 10a | List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect. | 8 |
| | 10b | List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information. | 8 |
| Study risk of bias assessment | 11 | Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process. | 8 |
| Effect measures | 12 | Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results. | 9 |
| Synthesis methods | 13a | Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)). | Supplement |
| | 13b | Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions. | 9 |
| | 13c | Describe any methods used to tabulate or visually display results of individual studies and syntheses. | 8-9 |
| | 13d | Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used. | 10 |
| | 13e | Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression). | 10 |
| | 13f | Describe any sensitivity analyses conducted to assess robustness of the synthesized results. | 10 |
| Reporting bias assessment | 14 | Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases). | 9 |
| Certainty assessment | 15 | Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome. | 9 |



PRISMA 2020 Checklist

| Section and Topic | Item # | Checklist item | Location where item is reported |
|--|--------|--|---------------------------------|
| RESULTS | | | |
| Study selection | 16a | Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram. | 11 |
| | 16b | Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded. | 11-12 |
| Study characteristics | 17 | Cite each included study and present its characteristics. | 8-9, Supplement |
| Risk of bias in studies | 18 | Present assessments of risk of bias for each included study. | Supplement |
| Results of individual studies | 19 | For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots. | Supplement |
| Results of syntheses | 20a | For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies. | Supplement |
| | 20b | Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect. | 11-13 |
| | 20c | Present results of all investigations of possible causes of heterogeneity among study results. | 11-13 |
| | 20d | Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results. | 11-13 |
| Reporting biases | 21 | Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed. | Supplement |
| Certainty of evidence | 22 | Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed. | Supplement |
| DISCUSSION | | | |
| Discussion | 23a | Provide a general interpretation of the results in the context of other evidence. | 14 |
| | 23b | Discuss any limitations of the evidence included in the review. | 16 |
| | 23c | Discuss any limitations of the review processes used. | 16 |
| | 23d | Discuss implications of the results for practice, policy, and future research. | 16 |
| OTHER INFORMATION | | | |
| Registration and protocol | 24a | Provide registration information for the review, including register name and registration number, or state that the review was not registered. | Supplement |
| | 24b | Indicate where the review protocol can be accessed, or state that a protocol was not prepared. | 7 |
| | 24c | Describe and explain any amendments to information provided at registration or in the protocol. | 7 |
| Support | 25 | Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review. | 17 |
| Competing interests | 26 | Declare any competing interests of review authors. | 17 |
| Availability of data, code and other materials | 27 | Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review. | Supplement |

BMJ Open

Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in healthcare workers: a meta-analysis of randomized clinical trials

| | |
|---------------------------------|--|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2022-065305.R3 |
| Article Type: | Original research |
| Date Submitted by the Author: | 29-May-2023 |
| Complete List of Authors: | Hong, Hwanhee; Duke University, Department of Biostatistics and Bioinformatics Friedland, Anne; UNC School of Medicine Hu, Mengyi; Duke University, Department of Biostatistics and Bioinformatics Anstrom, Kevin J.; University of North Carolina System, Collaborative Studies Coordinating Center Halabi, Susan; Duke University School of Medicine, Department of Biostatistics & Bioinformatics McKinnon, John; Henry Ford Hospital, Division of Infectious Diseases Amaravadi, Ravi; University of Pennsylvania Rojas-Serrano, Jorge; Instituto Nacional de Enfermedades Respiratorias, Interstitial Lung Disease and Rheumatology Units Abella, Benjamin; University of Pennsylvania Portillo-Vázquez, Angélica Margarita; Instituto Nacional de Enfermedades Respiratorias Woods, Christopher; Duke Clinical Research Institute Hernandez, Adrian F.; Duke University School of Medicine Boulware, David R.; University of Minnesota Twin Cities Naggie, Susanna; Duke Clinical Research Institute, Infectious Disease Rajasingham, Radha; University of Minnesota Twin Cities |
| Primary Subject Heading: | Infectious diseases |
| Secondary Subject Heading: | Infectious diseases, Evidence based practice |
| Keywords: | COVID-19, STATISTICS & RESEARCH METHODS, EPIDEMIOLOGY |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Safety and efficacy of hydroxychloroquine as prophylactic against COVID-19 in healthcare workers: a meta-analysis of randomized clinical trials

Hwanhee Hong, PhD (0000-0002-3736-6327), Anne Friedland, MD (0000-0002-9735-4838), Mengyi Hu, MB (0000-0003-1999-6672), Kevin J. Anstrom, PhD (0000-0001-6452-2172), Susan Halabi, PhD (0000-0003-4135-2777), John E. McKinnon, M.D., M.Sc (0000-0003-3916-4021), Ravi Amaravadi, MD (0000-0002-5768-2474), Jorge Rojas Serrano, MD, PhD (0000-0001-6980-7898), Benjamin S. Abella, MD MPhil (0000-0003-2521-0891), Angélica Margarita Portillo-Vázquez, MD, MSc (0000-0002-5032-4076), Christopher W. Woods, MD, MPH, Adrian Hernandez, MD, MHS (0000-0003-3387-9616), David R Boulware, MD MPH (0000-0002-4715-0060), Susanna Naggie, MD, MHS (0000-0001-7721-6975), Radha Rajasingham, MD (0000-0001-5531-0231)

Hwanhee Hong, Associate Professor, Department of Biostatistics and Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA

Anne Friedland, Assistant Professor, Department of Infectious Disease, University of North Carolina School of Medicine, 130 Mason Farm Rd, Chapel Hill, NC, 27514, USA

Mengyi Hu, Statistician, Department of Biostatistics and Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA

Kevin J. Anstrom, Director, Collaborative Studies Coordinating Center, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA

Susan Halabi, Professor, Department of Biostatistics and Bioinformatics, Duke University, 2424 Erwin Road, Suite 11088, Durham, NC, 27705, USA

1
2
3 23 John E. McKinnon, Senior Staff, Infectious Diseases, Henry Ford Health, Clara Ford Pavilion,
4
5 24 Suite 322, 2799 W. Grand Blvd., Detroit, MI 48202, USA
6
7 25 Ravi Amaravadi, Professor of Medicine, University of Pennsylvania , 852 BRB 2/3 421 Curie
8
9 26 Blvd., Philadelphia, PA 19063, USA
10
11
12 27 Jorge Rojas Serrano, Attending physician, Interstitial Lung Disease and Rheumatology Unit,
13
14 28 Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas, Calzada de Tlalpan
15
16 29 4502, Tlalpan, Ciudad de México, México. 14080
17
18
19 30 Benjamin S. Abella, Professor, Department of Emergency Medicine, University of Pennsylvania,
20
21 31 423 Guardian Drive Room 412, Philadelphia, PA 19104, USA
22
23 32 Angélica Margarita Portillo-Vázquez, Attending physician, Otolaryngology department, Instituto
24
25 33 Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas. Calzada de Tlalpan 4502,
26
27 34 Tlalpan, Ciudad de México, México. 14080
28
29
30 35 Christopher W. Woods, Professor of Medicine, Duke University, 310 Trent Drive, Durham, NC
31
32 36 27710, USA
33
34
35 37 Adrian Hernandez, Professor of Medicine, Duke Clinical Research Institute, Duke University
36
37 38 300 W. Morgan Street Durham, NC 27701, USA
38
39
40 39 David R Boulware, Professor, Division of Infectious Diseases & International Medicine,
41
42 40 Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis, MN, 55455,
43
44 41 USA
45
46
47 42 Susanna Naggie, Professor of Medicine, Duke Clinical Research Institute, 300 West Morgan
48
49 43 Street, 5th Floor, Durham, NC 27710, USA
50
51
52
53
54
55
56
57
58
59
60

44 Radha Rajasingham, Assistant Professor, Division of Infectious Diseases & International
45 Medicine, Department of Medicine, University of Minnesota, 689 23rd Ave SE, Minneapolis,
46 MN, 55455, USA

47
48 **Corresponding author:** Hwanhee Hong, Assistant Professor, Department of Biostatistics and
49 Bioinformatics, Duke University, 2424 Erwin Road, Ste 1105, Durham, NC, 27705, USA,
50 hwanhee.hong@duke.edu

51
52 Abstract word count: 225

53 Word count: 2473

54
55 **Key words:** SARS-CoV-2, COVID-19, hydroxychloroquine, prophylaxis, health care workers,
56 meta-analysis, clinical trials

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective: We studied the safety and efficacy of hydroxychloroquine (HCQ) as pre-exposure prophylaxis for COVID-19 in healthcare workers (HCWs), using a meta-analysis of randomized controlled trials.

Data Sources: PubMed, and EMBASE databases were searched to identify randomized trials studying HCQ.

Study Selection: Ten randomized controlled trials (RCTs) were identified (n=5,079 participants).

Data Extraction and Synthesis: The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis between HCQ and placebo using a Bayesian random-effects model. A *pre-hoc* statistical analysis plan was written, and the review protocol was registered at PROSPERO (CRD42021285093)

Main Outcomes: The primary efficacy outcome was polymerase chain reaction (PCR)-confirmed SARS-CoV-2 infection and the primary safety outcome was incidence of adverse events. The secondary outcome included clinically suspected SARS-CoV-2 infection.

Results: Compared with placebo, HCWs randomized to hydroxychloroquine (HCQ) had no significant difference in PCR-confirmed SARS-CoV-2 infection (odds ratio [OR] 0.92, 95% credible interval [CI]: 0.58, 1.37) or clinically suspected SARS-CoV-2 infection (OR 0.78, 95% CI: 0.57, 1.10), but significant difference in adverse events (OR 1.35, 95% CI: 1.03, 1.73).

Conclusions and Relevance: Our meta-analysis of ten RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in HCWs found that compared with placebo HCQ does not significantly reduce the risk of confirmed or clinically suspected SARS-CoV-2 infection, while HCQ significantly increases adverse events.

90

91 STRENGTHS AND LIMITATIONS OF THIS STUDY

- 92 • Bayesian meta-analysis models with random effects fitted the data.
- 93 • The ten trials included in the meta-analysis represent wide geographical locations
- 94 including US, Canada, Mexico, India, Spain, Bolivia, Venezuela, Peru, and Pakistan.
- 95 • The findings can be applied to healthcare workers but should not be generalized to a
- 96 broader population.

97

98 INTRODUCTION

99 Early during the SARS-CoV-2 pandemic, based on *in vitro* antiviral activity of both chloroquine
100 and hydroxychloroquine against SARS-CoV-2 [1-3], clinicians considered use of
101 hydroxychloroquine (HCQ) for treatment and prevention of SARS-CoV-2 infection and the
102 associated disease, COVID-19. While there are now published randomized controlled trials of
103 HCQ for the treatment of COVID-19 in the inpatient and outpatient setting [4, 5], there remains a
104 lack of adequately powered randomized controlled trials of HCQ for the pre-exposure
105 prophylaxis (PrEP) of SARS-CoV-2 infection. A number of COVID-19 clinical studies including
106 PrEP studies were planned early in the pandemic; however, several never opened to enrollment
107 and those that did open were closed early without reaching full accrual due to the rapidly
108 changing landscape of preventative therapies, including vaccines, and a significant shift in public
109 opinion of HCQ as a medical intervention for SARS-CoV-2 [6].

110 Vaccination access remains insufficient globally [7]. Specifically, in low-income
111 countries only 33% of healthcare workers are fully vaccinated. While high-income countries
112 have better coverage, overall 38% of countries did not achieve the milestone of 70% vaccination

coverage for healthcare workers by the end of 2021[8]. Thus, studying the pre-exposure prophylaxis potential for a drug with a known safety profile is crucial to protect people at high risk of exposures, such as healthcare workers (HCWs) [9, 10]. Two large randomized, placebo-controlled trials testing the safety and efficacy of HCQ as pre-exposure prophylaxis for COVID-19 in HCWs [11] [12], showed potential for a modest benefit of HCQ but were both underpowered, if a modest effect exists. More trials [13-15] studying HCQ as pre-exposure prophylaxis of COVID-19 in HCWs have been published with similar limitations.

To address the most common limitation, inadequate power to show a modest effect, we conducted a formal meta-analysis of pre-exposure prophylactic HCQ studies in HCWs. We conducted a systematic search for clinical trials of pre-exposure prophylactic use of HCQ against infection of SARS-CoV-2 in HCWs, thoroughly compared similarities and differences in characteristics of the identified studies and performed a Bayesian meta-analysis to combine results of the trials.

METHODS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used in this systematic review and meta-analysis[16]. A statistical analysis plan was written in advance and the review protocol was registered at PROSPERO (CRD42021285093).

Search strategy and information sources

We searched PubMed/Medline and Ovid/Embase databases from database inception through the final search date March 14, 2023. We used keywords related to COVID-19, HCQ, and randomized controlled trials. The full search strategies are provided in eTable 1.

136

137 **Eligibility criteria and study selection**

138 The eligibility criteria included phase II or phase III randomized controlled trials (RCTs) of
139 hydroxychloroquine for use as pre-exposure prophylaxis in HCWs with moderate to high risk of
140 exposure. We excluded observational studies, crossover trials, studies where the method of
141 allocation to treatment was not truly random, duplicate studies, and non-original data studies. No
142 language, publication date, or publication status restrictions were applied. References of prior
143 systematic reviews and meta-analyses were also screened for related studies. Study selection
144 involved screening of titles and abstracts followed by full-text evaluation of possible eligible
145 studies.

146

147 **Data collection process**

148 Each of the selected studies were independently reviewed by two reviewers (AF, MH, or HH).
149 We extracted data on the study design, baseline characteristics, interventions, and outcomes. Any
150 disagreements of collected information between reviews were reconciled through discussion by
151 all three reviewers.

152

153 **Outcome measures**

154 The primary efficacy outcome for the meta-analysis was laboratory confirmed SARS-CoV-2
155 infection by polymerase chain reaction (PCR) test and the primary safety outcome was incidence
156 of adverse events (Table 1). The secondary efficacy outcome was suspected or probable SARS-
157 CoV-2 infection. Included studies had the following outcome definitions: (1) laboratory
158 confirmed SARS-CoV-2 infection defined as COVID-19 like symptoms and positive SARS-

CoV-2 PCR and (2) suspected or probable SARS-CoV-2 infection defined as COVID-19 like symptoms but lack of confirmatory PCR testing.

Table 1. Treatment strategies, adherence, trial-defined primary outcome, and study duration for trials included in the meta-analysis

| | Trial-defined primary outcome | Study duration | Treatment group | Randomized treatment assignment | Randomized sample size |
|--|--|---------------------|------------------|--|------------------------|
| Naggie et al.[13] (HERO-HCQ) | Confirmed (by NP swab PCR) or suspected COVID-19 infection through 30 days | 60 days | HCQ | HCQ 600 mg BID loading dose for Day 1, followed by 400 mg QD for 29 days | 683 |
| | | | Control | Placebo | 676 |
| Abella et al.[11] (PATCH) | COVID-19 infection as determined by positive NP swab over 8 weeks | 56 days (8 weeks) | HCQ | HCQ 600mg daily for 60 days | 64 |
| | | | Control | Placebo | 61 |
| Rajasingham et al.[12] (MN-COVID-PREP) | COVID-19 free survival time by lab confirmed or probable illness | 84 days (12 weeks) | HCQ ^a | HCQ loading doses (400 mg twice 6-8hrs apart), followed by 400 mg once weekly or 400 mg twice weekly for 84 days | 989 |
| | | | Control | Placebo | 494 |
| Rojas-Serrano et al.[14] | Time to symptomatic respiratory infection with a positive COVID RT PCR over 60 days | 60 days | HCQ | HCQ 200 mg daily for 60 days | 62 |
| | | | Control | Placebo | 65 |
| McKinnon et al.[15] (WHIP) | Lab confirmed cases of COVID-19 determined by either IgM and IgG serology in blood sample or RT-PCR test results | 56 days (8 weeks) | HCQ ^a | HCQ 400 mg loading dose for Day 1, followed by 200 mg daily or 400 mg weekly on the same day of each week for 56 days | 387 |
| | Confirmed new cases of COVID-19 | | Control | Placebo | 191 |
| Vijayaraghavan et al.[17] | Lab confirmed SARS-CoV-2 infection by PCR or presence of antibodies | 180 days (6 months) | HCQ | HCQ 400 mg twice on the day of enrollment, followed by 400 mg once a week for a total of 12 weeks plus personal protective equipment (PPE) | 213 |
| | | | Control | PPE | 203 |
| Polo et al.[18] (EPICOS) | Lab confirmed symptomatic COVID-19 by PCR | 84 days (12 weeks) | HCQ ^b | HCQ 200 mg once daily | 231 |
| | | | Control | Placebo | 223 |
| Llanos-Cuentas et al.[19] | COVID-19 cases confirmed by PCR or serological test | 28 days (4 weeks) | HCQ | HCQ loading dose of 600 mg on the first day, followed by 400 mg every other day plus PPE | 36 |
| | | | Control | PPE | 32 |
| Grau-Pujol et al.[20] | COVID-19 confirmed cases with | 180 days (6 months) | HCQ | HCQ 400 mg daily for the four consecutive | 142 |

| PCR test | | | days, followed by 400 mg weekly | | |
|-----------------|--|--------------------|---------------------------------|---|-----|
| | | | Control | Placebo | 127 |
| Syed et al.[17] | COVID-19-free survival (COVID-19 confirmed by PCR) | 84 days (12 weeks) | HCQ ^a | HCQ 400 mg twice for Day 1, followed by 400 mg weekly or HCQ 400 mg once every 3 weeks or HCQ 200 mg once every 3 weeks | 154 |
| | | | Control | Placebo | 46 |

HCQ=Hydroxychloroquine

^a More than one HCQ groups with different doses are lumped.

^b The Polo et al. study randomized participants to four treatment groups, and the HCQ and control groups are used in our meta-analysis.

Treatment assignment

Our meta-analysis did not study HCQ dosing specific effects. For studies randomizing participants to more than one HCQ arm with different doses, all HCQ arms were merged and considered as a single HCQ arm. Such studies include the Rajasingham et al., McKinnon et al. and Syed et al. studies.

Risk of bias and certainty of evidence assessment

Two independent reviewers (AF, HH) assessed the risk of bias (low, intermediate, high) of the included studies using the Cochrane's Collaboration tool [21] (eTable 2). We assessed the certainty of evidence using the Grades of Recommendation Assessment, Development and Evaluation (GRADE) approach [22].

Statistical analysis

Bayesian logistic regression meta-analysis models under two assumptions (fixed effect and random effects) were fitted to estimate the odds ratio of having an outcome between hydroxychloroquine and placebo [23]. The fixed effect model assumes that the odds ratio is constant across studies, while the random effects model accounts for heterogeneity in the odds

1
2
3 187 ratios across studies. To assess and compare the goodness-of-fit of the fitted fixed and random
4
5 188 effects models, we calculated the Watanabe-Akaike information criterion [24]. In the Bayesian
6
7 189 models, we assigned non-informative prior distributions as no prior information was available.
8
9
10 190 The odds ratios and the associated 95% credible intervals were estimated using Markov chain
11
12 191 Monte Carlo (MCMC) algorithms. In addition, we calculated Bayesian posterior probabilities of
13
14 192 the odds ratio smaller than 1 or 0.5 for the primary efficacy outcome, and greater than 2 for the
15
16 193 safety outcome [25]. The standard deviation of the random effects and I^2 [26] were estimated to
17
18 194 quantify the between-study heterogeneity, where small values of both metrics indicate slight
19
20 195 heterogeneity. To identify publication bias, we plotted and assessed funnel plots for their
21
22 196 symmetry, and conducted the Egger's test[27]. All Bayesian meta-analyses were conducted
23
24 197 using the `rstan` package (version 2.21.2)[28] in R 4.0.2 [29]. We used two parallel chains,
25
26 198 where each chain consists of 50,000 samples after a 25,000-sample burn-in. We checked
27
28 199 convergence of the MCMC chains for all model parameters using trace plots and Gelman-Rubin
29
30 200 diagnostic statistics [30].
31
32
33
34
35
36
37

38 202 **Patient and public involvement**

39
40 203 No patient involved.
41
42
43

44 204
45 205 **RESULTS**

46
47 206 **Search results**

48
49 207 Our database search resulted in 350 unique studies after excluding duplicates. Of those, 339
50
51 208 studies were screened out due to irrelevance based on title and abstract screening. Eleven studies
52
53 209 were assessed in full-text for eligibility (Figure 1). Of those, one trial was excluded from the
54
55
56
57
58
59
60

meta-analysis because it studied with non-healthcare worker populations. As a result, a total of ten studies in a population consisting of HCWs were identified (Table 1).

Study and patient characteristics

Study design, population, treatment strategies, and key characteristics are presented in Table 1 and eTable 3. A total of 5,079 randomized participants (2,961 randomized to HCQ) from the 10 studies were included in the meta-analysis. The ten studies defined HCWs broadly and included first responders (emergency medical services, fire, and police). The follow-up duration of the 10 studies ranged from 28 days to 180 days. The HCQ dosing scheme varied across studies, including daily dosing ranging from 200 to 600mg daily with or without a loading dose and once or twice weekly or once every three weeks dosing. The duration of therapy also varied across studies (Table 1). The trial-specific definitions of primary outcome and adverse events are comparable across trials (Table 1, eTable 4).

Baseline characteristics by randomized treatment assignment are reported (eTable 5). The average age ranged between 31 and 45. The aggregate proportion of women within each study varied across the 10 trials, with a range from 44% to 69%. In addition, the Abella et al. and Rojas-Serrano et al. studies had smaller sample size compared with the other three studies and showed a difference in female ratio between placebo and HCQ groups. In the Naggie et al., Abella et al., Rajasingham et al., and McKinnon et al., studies, over 80% of study participants were white. The Abella et al. and Rajasingham et al. studies had high proportions of HCWs working in an emergency department (56% and 41%, respectively) and the Abella et al. study had a high proportion of nurses (67%).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

Several studies reported treatment adherence assessed by two methods: self-reported adherence and/or pill count at the end of the study. The Rajasingham et al. study additionally conducted remote blood sampling to verify HCQ concentrations in a subset. Adherence varied significantly across the studies, with a low proportion of approximately 52% in the Rojas-Serrano et al. study and 97-98% in the Abella et al. study.

Results of meta-analysis

Overall, 3.4% (171/5039) developed PCR-confirmed SARS-CoV-2 infection and 5.6% (230/4087) developed suspected COVID-19 that was not laboratory confirmed. Since the goodness-of-fit assessment using Watanabe-Akaike information criterion concluded that the random effects meta-analysis model was as good as or better than the fixed effect meta-analysis model for all outcomes, we reported the results under the random effects model. Compared with placebo, HCWs randomized to HCQ had numerically lower rate of PCR-confirmed SARS-CoV-2 infection cases (odds ratio [OR] 0.92, 95% credible interval [CI]: 0.58, 1.37; GRADE score: moderate certainty), and suspected or probable SARS-CoV-2 infection cases (OR 0.78, 95% CI: 0.57, 1.10; GRADE score: moderate certainty). None of these odds ratios were statistically significant. Participants treated with HCQ had a numerically higher rate of adverse events (OR 1.35, 95% CI: 1.03, 1.73; GRADE score: moderate certainty) with statistical significance (Figure 2). The outcome data used in our analyses are presented in eTable 6. The summary of GRADE score assessment is provided in eTable 7.

The Bayesian posterior probabilities of the odds ratio less than 1 for the confirmed SARS-CoV-2 infection outcome (i.e., the probability of HCQ favoring over placebo) was 0.67, while the posterior probability of odds ratio less than 0.5 (i.e., the probability that the odds of having a confirmed SARS-CoV-2 infection outcome in HCQ is less than a half of the odds in placebo) was 0.009. The posterior probability of the odds ratio greater than 2 for the adverse event outcome (i.e., the probability that the odds of having an adverse event in HCQ is greater than twice of the odds in placebo) was 0.004.

Our meta-analysis showed little or moderate variability of effect estimates across studies with I^2 value of 0%, 0%, and 43%, and the estimated standard deviation of the random effects of 0.39, 0.26, and 0.45 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively. Funnel plots (eFigure) showed no indication of publication bias and the associated Egger's test results supported that the funnel plots were not asymmetry with p-values of 0.308, 0.305, and 0.794 for the confirmed SARS-CoV-2 infection, suspected SARS-CoV-2 infection, and adverse event outcomes, respectively.

DISCUSSION

Understanding the pre-exposure prophylactic effect of HCQ against COVID-19 remains relevant, as its use continues, particularly in the international setting [31, 32]. Our meta-analysis of the ten RCTs investigating the safety and efficacy of HCQ as pre-exposure prophylaxis in 5,079 HCWs found that HCQ did not have a statistical association with fewer confirmed or suspected/probable SARS-CoV-2 infection cases compared with placebo. The geographical locations of the 10 trials included in the meta-analysis are US, Canada, Mexico, India, Spain,

1
2
3 278 Bolivia, Venezuela, Peru, and Pakistan (eTable 3). While the odds ratios of most studies favor
4
5 279 HCQ, the credible intervals remain wide suggesting low certainty in the true point estimate. Two
6
7
8 280 studies including the Llanos-Cuentas et al. study conducted in Peru and the Syed et al. study
9
10 281 conducted in Pakistan showed odds ratios favoring placebo, though the credible intervals remain
11
12 282 wide. Furthermore, in this population, COVID-19 events rates were low, particularly for the
13
14 283 most relevant PCR-confirmed infection outcome. The low event rate raises further concern for
15
16 284 the uncertainty of these outcomes. Thus, if there is a minimal effect, the absolute benefit would
17
18 285 be low. To gain more certainty, a very large study would need to be done and this is difficult to
19
20 286 support now due to availability of highly effective vaccines. The safety profile of HCQ in the
21
22 287 outpatient setting is well understood [33]. In these outpatient studies there was statistically
23
24 288 significant difference in adverse events in the HCQ versus the placebo arm, indicating that HCQ
25
26 289 is less safe than placebo.
27
28
29
30
31 290
32
33 291 Our findings can be applied to HCWs but should not be generalized to a broader population. Our
34
35 292 systematic search found only one published RCT of pre-exposure prophylaxis for non-healthcare
36
37 293 worker populations and the study were excluded from our meta-analysis. This study was
38
39 294 conducted in Singapore [34] and showed a significant reduction in the risk of COVID-19
40
41 295 infection in the HCQ arm when compared with the comparator arm, vitamin C. However, this
42
43 296 study showed moderate risk of bias as it used an open-label cluster-randomization design, the
44
45 297 Institutional Review Board excluded higher risk persons from the hydroxychloroquine arm only,
46
47 298 and the participants may not be representative of a general population due to the communal
48
49 299 living environment.
50
51
52
53
54 300
55
56
57
58
59
60

1
2
3 301 A Bayesian meta-analysis approach was used to fit the data. The Bayesian meta-analysis
4
5 302 approach has several advantages. First, its flexibility and the MCMC sampling methods to
6
7 303 estimate posterior distributions provide probability-based quantities (e.g., posterior probability of
8
9 304 an odds ratio smaller than 0.5) that complement typical meta-analysis results (e.g., odds ratios
10
11 305 and the associated credible intervals) and help decision making [35]. Second, the Bayesian meta-
12
13 306 analysis model with random effects estimates the between-study variability better than the
14
15 307 frequentist counterparts [36]. Third, when it comes to with binary outcomes, the Bayesian
16
17 308 approach handles rare events better than the frequentist counterparts [23].
18
19
20
21
22 309

23
24 310 A recently published meta-analysis by García-Albéniz et al. [37] investigated pre-exposure
25
26 311 (seven RCTs included) and post-exposure (four RCTs included) prophylactic effects of HCQ,
27
28 312 but not limited to the HCW population. They found significant pre-exposure prophylactic effects
29
30 313 of HCQ on SARS-CoV-2 infection, different from ours. The seven pre-exposure prophylaxis
31
32 314 RCTs included in the García-Albéniz et al. meta-analysis consisted of six RCTs that were in our
33
34 315 meta-analysis and the aforementioned Singapore study that was excluded from our meta-
35
36 316 analysis. Our meta-analysis provides the most up-to-date, systematic, and comprehensive
37
38 317 evidence about prophylactic effects of HCQ focusing on the HCW population.
39
40
41
42 318

43
44 319 Although a meta-analysis allows for combining evidence from multiple studies in a principled
45
46 320 way, our meta-analysis has limitations. First, our analysis did not evaluate effects of different
47
48 321 HCQ doses and combined multiple HCQ arms using different doses in three studies. The RCTs
49
50 322 included in our meta-analysis studied varying dosing schemes and a meta-analysis using
51
52 323 aggregate-level data is not a sufficient source to study dosing effects. Second, detailed subgroup
53
54
55
56
57
58
59
60

1
2
3 324 analyses were not conducted due to limited information. Individual-level data are required to
4
5 325 study both dosing and subgroup effects.
6
7 326
8
9
10 327 Our meta-analysis of ten RCTs investigating safety and efficacy of HCQ as pre-exposure
11
12 328 prophylaxis in HCWs provides the most up-to-date evidence on HCQ. Although most individual
13
14 329 trials were underpowered and showed null data, integrating the results systematically via meta-
15
16 330 analysis contributes to the scientific literature and provides certain answers to the question. We
17
18 331 found that HCQ does not reduce the risk of confirmed or probable SARS-CoV-2 infection, but
19
20 332 increase risk of adverse events compared with placebo. Hydroxychloroquine should not be used
21
22 333 for pre-exposure prophylaxis in the HCW population.
23
24
25
26
27

28 335 **Contributors**

29
30
31 336 All authors fulfill the ICMJE criteria for authorship. HH, SN, RR, and KJA designed the study.
32
33 337 HH, AF, and MH collected and analyzed the data. HH, SN, and RR wrote the manuscript. SH
34
35 338 and KJA provided statistical review and AF, JEM, RA, JRS, BSA, AMPV, CWW, AH and DRB
36
37 339 provided clinical review. All authors approved and decided to submit the paper for publication.
38
39
40 340 Hwanhee Hong – HH
41
42 341 Anne Friedland – AF
43
44 342 Mengyi Hu – MH
45
46 343 Kevin J. Anstrom – KJA
47
48 344 Susan Halabi – SH
49
50 345 John E. McKinnon – JEM
51
52 346 Ravi Amaravadi – RA
53
54
55
56
57
58
59
60

347 Jorge Rojas Serrano – JRS

348 Benjamin S. Abella – BSA

349 Angélica Margarita Portillo-Vázquez – AMPV

350 Christopher W. Woods – CWW

351 Adrian Hernandez – AH

352 David R Boulware – DRB

353 Susanna Naggie – SN

354 Radha Rajasingham – RR

355 **Funding**

356 This study is funded by the Patient Centered Outcomes Research Institute (PCORI), Contract
357 Number COVID-19-2020-001. The funder had no role in the design, conduct, analysis, or
358 reporting of this study.

359 **Competing interests**

360 All authors except Dr. Abella reported no financial relationship with commercial interest. Dr.
361 Abella have received NIH funds for COVID-19 related research, and holds equity in VOC
362 Health, a start-up company that is developing novel covid testing.

363 **Ethics Approval**

364 Ethics approval was not required because this study used publicly available aggregate data that
365 were not involved with patients' information or prospective data collection.

366 **Data sharing statement**

367 The data are presented in eTable 6.

369 **REFERENCES**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Kalil, A.C., *Treating COVID-19—off-label drug use, compassionate use, and randomized clinical trials during pandemics*. JAMA, 2020. **323**(19): p. 1897-1898.

2. McCreary, E.K., J.M. Pogue, and o.b.o.t.S.o.I.D. Pharmacists, *Coronavirus Disease 2019 Treatment: A Review of Early and Emerging Options*. Open Forum Infectious Diseases, 2020. **7**(4).

3. Wang, M., et al., *Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro*. Cell research, 2020. **30**(3): p. 269-271.

4. RECOVERY Collaborative Group, *Effect of hydroxychloroquine in hospitalized patients with Covid-19*. New England Journal of Medicine, 2020. **383**(21): p. 2030-2040.

5. Skipper, C.P., et al., *Hydroxychloroquine in nonhospitalized adults with early COVID-19: a randomized trial*. Annals of internal medicine, 2020. **173**(8): p. 623-631.

6. Halabi, S., et al., *Landscape of coronavirus disease 2019 clinical trials: New frontiers and challenges*. Clinical Trials, 2022: p. 17407745221105106.

7. Padma, T., *COVID vaccines to reach poorest countries in 2023—despite recent pledges*. Nature, 2021. **595**(7867): p. 342-343.

8. Nabaggala, M.S., et al., *The global inequity in COVID-19 vaccination coverage among health and care workers*. International Journal for Equity in Health, 2022. **21**(3): p. 147.

9. World Health Organization. *Prevention, identification and management of health worker infection in the context of COVID-19*. 2020 [cited 2022 May 13th]; Available from: <https://www.who.int/publications/i/item/10665-336265>

10. The United Kingdom Office for National Statistics. *Coronavirus (COVID-19) infections in the community in England: May 2021*. 2021 [cited 2022 May 13th]; Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsinthecommunityinengland/characteristicsofpeopletestingpositiveforcovid19incountriesoftheuk20may2021#percentage-testing-positive-for-covid-19-by-patient-facing-and-non-patient-facing-job-roles-uk>.

11. Abella, B.S., et al., *Efficacy and safety of hydroxychloroquine vs placebo for pre-exposure SARS-CoV-2 prophylaxis among health care workers: a randomized clinical trial*. JAMA internal medicine, 2021. **181**(2): p. 195-202.

12. Rajasingham, R., et al., *Hydroxychloroquine as Pre-exposure Prophylaxis for Coronavirus Disease 2019 (COVID-19) in Healthcare Workers: A Randomized Trial*. Clinical Infectious Diseases, 2020. **72**(11): p. e835-e843.

13. Naggie, S., et al., *Hydroxychloroquine for pre-exposure prophylaxis of COVID-19 in health care workers: a randomized, multicenter, placebo-controlled trial Healthcare Worker Exposure Response and Outcomes of Hydroxychloroquine (HERO-HCQ)*. International Journal of Infectious Diseases, 2023. **129**: p. 40-48.

14. Rojas-Serrano, J., et al., *Hydroxychloroquine for prophylaxis of COVID-19 in health workers: A randomized clinical trial*. PLoS One, 2022. **17**(2): p. e0261980.

15. McKinnon, J.E., et al., *Safety and tolerability of hydroxychloroquine in health care workers and first responders for the prevention of COVID-19: WHIP COVID-19 Study*. International Journal of Infectious Diseases, 2022. **116**: p. 167-173.

16. Hutton, B., et al., *The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations*. Annals of internal medicine, 2015. **162**(11): p. 777-784.

17. Tirupakuzhi Vijayaraghavan, B.K., et al., *Hydroxychloroquine plus personal protective equipment versus personal protective equipment alone for the prevention of laboratory-confirmed COVID-19 infections among healthcare workers: a multicentre, parallel-group randomised controlled trial from India*. BMJ Open, 2022. **12**(6): p. e059540.
18. Polo, R., et al., *Daily tenofovir disoproxil fumarate/emtricitabine and hydroxychloroquine for pre-exposure prophylaxis of COVID-19: a double-blind placebo-controlled randomized trial in healthcare workers*. Clinical Microbiology and Infection, 2023. **29**(1): p. 85-93.
19. Llanos-Cuentas, A., et al., *Hydroxychloroquine to prevent SARS-CoV-2 infection among healthcare workers: early termination of a phase 3, randomised, open-label, controlled clinical trial*. BMC Research Notes, 2023. **16**(1): p. 22.
20. Grau-Pujol, B., et al., *Pre-exposure prophylaxis with hydroxychloroquine for COVID-19: a double-blind, placebo-controlled randomized clinical trial*. Trials, 2021. **22**(1): p. 808.
21. Sterne, J.A., et al., *RoB 2: a revised tool for assessing risk of bias in randomised trials*. bmj, 2019. **366**.
22. Puhan, M.A., et al., *A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis*. Bmj, 2014. **349**.
23. Hong, H., C. Wang, and G.L. Rosner, *Meta-analysis of rare adverse events in randomized clinical trials: Bayesian and frequentist methods*. Clinical Trials, 2021. **18**(1): p. 3-16.
24. Watanabe, S. and M. Opper, *Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory*. Journal of machine learning research, 2010. **11**(12).
25. Ferreira, D., et al., *Bayesian predictive probabilities: a good way to monitor clinical trials*. British journal of anaesthesia, 2021. **126**(2): p. 550-555.
26. Higgins, J.P. and S.G. Thompson, *Quantifying heterogeneity in a meta-analysis*. Statistics in medicine, 2002. **21**(11): p. 1539-1558.
27. Egger, M., et al., *Bias in meta-analysis detected by a simple, graphical test*. Bmj, 1997. **315**(7109): p. 629-634.
28. Stan Development Team, *RStan: the R interface to Stan*. R package version, 2020. **2.21.2**.
29. R Core Team, *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2021.
30. Gelman, A. and D.B. Rubin, *Inference from iterative simulation using multiple sequences*. Statistical science, 1992. **7**(4): p. 457-472.
31. Infante, M., et al., *Hydroxychloroquine in the COVID-19 pandemic era: in pursuit of a rational use for prophylaxis of SARS-CoV-2 infection*. Expert review of anti-infective therapy, 2021. **19**(1): p. 5-16.
32. *Revised advisory on the use of hydroxychloroquine (HCQ) as prophylaxis for SARS-CoV-2 infection (in supersession of previous advisory dated 23rd March. 2020)*. 2022; Available from: https://www.icmr.gov.in/pdf/covid/techdoc/V5_Revised_advisory_on_the_use_of_HCQ_SARS_CoV2_infection.pdf.
33. Lofgren, S.M., et al. *Safety of hydroxychloroquine among outpatient clinical trial participants for COVID-19*. in *Open forum infectious diseases*. 2020. Oxford University Press US.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

34. Seet, R.C.S., et al., *Positive impact of oral hydroxychloroquine and povidone-iodine throat spray for COVID-19 prophylaxis: An open-label randomized trial*. International Journal of Infectious Diseases, 2021. **106**: p. 314-322.

35. Hong, H., et al., *A Bayesian missing data framework for generalized multiple outcome mixed treatment comparisons*. Research synthesis methods, 2016. **7**(1): p. 6-22.

36. Hong, H., et al., *Comparing Bayesian and frequentist approaches for multiple outcome mixed treatment comparisons*. Medical Decision Making, 2013. **33**(5): p. 702-714.

37. García-Albéniz, X., et al., *Systematic review and meta-analysis of randomized trials of hydroxychloroquine for the prevention of COVID-19*. European Journal of Epidemiology, 2022. **37**(8): p. 789-796.

Figure Legends

Figure 1. Flowchart of literature review

Figure 2. Forest plots of the meta-analysis results showing the number of events (y), sample size (n), posterior median of odds ratios, and the associated 95% credible intervals comparing HCQ versus placebo for (a) lab-confirmed positive COVID-19, (b) suspected COVID-19, and (c) adverse events.

For peer review only

1 Identification
2
3
4
5

6
7

8
9

10
11

12
13

14
15

16
17

18
19

20
21

22
23

24
25

26
27

28
29

30
31

32
33

34
35

36
37

38
39


40
41


42
43

44
45


46
47

48






(b) Suspected COVID-19



(c) Adverse events



Supplementary Materials

CONTENTS

- eTable 1. Search code
- eTable 2. Risk of bias
- eTable 3. Characteristics of included trials
- eTable 4. Definition of adverse events
- eTable 5. Baseline characteristics
- eTable 6. Results of outcome measures in each study
- eFigure. Funnel plots for the three outcomes
- eTable 7. GRADE summary of findings table



















































eTable 1. Search code that was used to identify publications as of March 14, 2023**PubMed search**

| | |
|----|--|
| #1 | covid[Title] OR coronavirus[Title] OR sars-cov-2[Title] |
| #2 | hydroxychloroquine[Title] |
| #3 | randomized[Title/Abstract] OR randomized[Title/Abstract] |
| #4 | #1 AND #2 AND #3 |

Embase search

| | |
|----|---|
| #1 | covid:ti OR coronavirus:ti OR 'sars cov 2':ti |
| #2 | hydroxychloroquine:ti |
| #3 | randomized:ab,ti OR randomised:ab,ti |
| #4 | #1 AND #2 AND #3 |

eTable 2. Risk of bias for trials included in the meta-analysis using the Cochrane risk assessment tool. Green circle is for low risk and yellow circle is for some concerns

| | Selection bias (Randomization process) | Performance bias (Deviations from the intended interventions) | Attrition bias ¹ (Missing outcome data) | Reporting bias (Measurement of the outcome) | Other sources of bias (Selection of the reported result) |
|---------------------------------------|---|---|---|---|---|
| Naggie et al. (HERO-HCQ) |  |  |  |  |  |
| Abella et al. (PATCH) |  |  |  |  |  |
| Rajasingham et al. (MN-COVID-PREP) |  |  |  |  |  |
| Rojas-Serrano et al. |  |  |  |  |  |
| McKinnon et al. (WHIP) |  |  |  |  |  |
| Vijayaraghavan et al. |  |  |  |  |  |
| Polo et al. (EPICOS) |  |  |  |  |  |
| Llanos-Cuentas et al. |  |  |  |  |  |
| Grau-Pujol et al. |  |  |  |  |  |
| Syed et al. |  |  |  |  |  |

¹ The Rojas-Serrano et al. study reported minimal loss to follow-up (<10%). The Rojas-Serrano et al. study reported 18% (25/130) lost to follow-up and additional 12% (16/130) discontinued the intervention.

eTable 3. Characteristics of trials included in the meta-analysis

| | Naggie et al. (HERO-HCQ) | Abella et al. (PATCH) | Rajasingham et al. (MN-COVID-PREP) | Rojas-Serrano et al. | McKinnon et al. (WHIP) |
|---|--|--|--|---|--|
| N (randomization) | 1360 | 132 | 1496 | 130 | 624 |
| Study start date ¹ | 4/22/2020 | 4/9/2020 | 4/6/2020 | 4/21/2020 | 4/10/2020 |
| Study completion date ² | 1/9/2021 | 11/13/2020 | 7/13/2020 | 3/31/2021 | 12/14/2020 |
| Occupation | HCWs at risk of COVID exposure through work in the ICU, emergency department, emergency services, respiratory services or COVID unit | HCWs (Physicians, nurses, certified nursing assistants, emergency technicians, respiratory therapists) eligible working >20 hrs/week | HCWs (physicians, nurses, emergency medical technicians) with direct contact with COVID patients including emergency department and ICU setting, first responders and performing aerosol generating procedures | HCWs (nurses, nursing aids, cleaning staff, orderlies, respiratory therapists and physicians) taking care of hospitalized patients with COVID | HCW, first responders and correlational/law officers, nursing home workers, medical students, public transit workers, household family members of HCW in Michigan and Ohio |
| Sites | 34 sites across the US | 2 tertiary urban hospitals | Multiple sites nationwide across US and Canada | Single site (National Institute of Respiratory Diseases of Mexico) | Multiple sites at Michigan in the US |
| Randomization | Yes (Phase III) | Yes (Phase II) | Yes (Phase III) | Yes (Phase III) | Yes (Phase III) |
| Trial type | Double-blinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | No weight requirement | <40kg excluded | <50kg excluded | N/A |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Excluded | Excluded | Excluded | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Included | Exclude |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Included | |
| Significant renal disease (stage IV or greater) | Excluded | Included | Excluded | Excluded | Exclude |
| Pregnant/breastfeeding | Included | Excluded | Included in US, Excluded in Canada | Excluded | Exclude |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Excluded | Included | Exclude |
| Use of other medications with significant drug interactions | Included | Excluded | Excluded | Included | N/A |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine or azithromycin) | Any treatment for COVID-19 within 14 days excluded | Current use of HCQ or chloroquine excluded | HCQ or chloroquine within 30 days excluded | Chronic use of HCQ included |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | N/A | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Excluded | Excluded if symptoms within 2 weeks unless negative COVID test | Excluded | Excluded | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | N/A |
| Positive COVID serology | Included | Included | N/A | Included | N/A |
| Analysis | Modified intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat | Intention-to-treat |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

| | Vijayaraghavan et al. | Polo et al. (EPICOS) | Llanos-Cuentas et al. | Grau-Pujol et al. | Syed et al. |
|---|---|--|--|--|--|
| N (randomization) | 416 | 454 | 68 | 269 | 200 |
| Study start date ¹ | 6/29/2020 | 4/2020 Spain, 10/2020 Bolivia, 3/2021 Venezuela | June, 2020 | 4/4/2020 | 5/1/2020 |
| Study completion date ² | 2/4/2021 | 5/30/2021 | November, 2020 | Study halted a 1 month analysis | Not reported |
| Occupation | HCWs in an environment with exposure to COVID-19 (physicians, nurses, allied health workers and ancillary health workers) | HCWs (physicians, nurses, medical students, other workers with and without direct patient contact) | HCWs (physicians, nursing staff, technical staff and nursing assistants involved in care of COVID-19 patients) | HCWs (physicians, nurses, nurse assistants and administrators working at least 3 days a week in the trial hospitals) | HCWs at risk of COVID-19 exposure including physicians, nurses, first responders, those performing aerosol generating procedures or working in the emergency department, ICU, and general medicine wards |
| Sites | 9 hospitals across India | Multiple sites across Spain, Venezuela and Bolivia | 4 public hospitals across the Lima metropolitan area | 3 hospitals in Barcelona, Spain | Single hospital in Pakistan |
| Randomization | Yes | Yes | Yes (Phase III) | Yes | Yes (Phase II) |
| Trial type | Unblinded | Double-blinded | Double-blinded | Double-blinded | Double-blinded |
| Eligibility criteria | | | | | |
| Age | >18 | >18-70 | >18 | >18 | >18 |
| Sex | All | All | All | All | All |
| Weight | No weight requirement | <40kg excluded | No weight requirement | No weight requirement | <40 kg |
| Health conditions | | | | | |
| Allergy or hypersensitivity to HCQ | Excluded | Excluded | Excluded | Excluded | Excluded |
| G6PD deficiency | Included | Included | Excluded | Included | Exclude |
| H/o retinal disease | Excluded | Excluded | Excluded | Excluded | Excluded |
| History of significant cardiac disease or Qtc prolongation | Excluded | Excluded | Excluded | Excluded | Excluded |
| Significant renal disease (stage IV or greater) | Included | Excluded | Excluded | Excluded | Excluded |
| Pregnant/breastfeeding | Excluded | Excluded | Included | Excluded | Excluded |
| Medication | | | | | |
| Qtc prolonging medications | Excluded | Excluded | Included | Excluded | Excluded |
| Use of other medications with significant drug interactions | Excluded | Included | Included | Excluded | Excluded |
| HCQ or other COVID treatments | Excluded (hydroxychloroquine, chloroquine azithromycin) | Any medication as prophylaxis against COVID-19 after 3/1/21 | Use of hydroxychloroquine, chloroquine or azithromycin in the last 30 days excluded | Treatment with chloroquine or hydroxychloroquine within the last 1 month | Those already taking hydroxychloroquine were excluded |
| COVID-19 related criteria | | | | | |
| Active or prior COVID | Excluded | Excluded | Excluded | Excluded | Excluded |
| Fevers, cough, SOB | Not specified in exclusion criteria | Excluded | Not specified in exclusion criteria | Not specified in exclusion criteria | Excluded |
| Positive COVID PCR | Excluded | Excluded | Excluded | Excluded | Excluded |
| Positive COVID serology | N/A | N/A | N/A | Excluded | Excluded |
| Analysis | Intention-to-treat | Not reported | Intention-to-treat | Intention-to-treat | Not reported |

HCW=Healthcare workers; ICU=Intensive care unit; ¹ Date when first participant was enrolled; ² Date when final data were collected for the last participant

eTable 4. Definition of adverse events

| Trial | AE definition |
|---|---|
| Naggie et al. (HERO-HCQ) | Adverse events include general disorders and administration site conditions, psychiatric disorders, skin and subcutaneous tissue disorders, cardiac disorders, infections and infestations, nervous system disorders, gastrointestinal disorders, investigations (electrocardiogram QT prolonged and heart rate increased), ear and labyrinth disorders, renal and urinary disorders, and respiratory, thoracic and mediastinal disorders. |
| Abella et al. (PATCH) | Adverse events include abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. |
| Rajasingham et al. (MN-COVID-PREP) | Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and others. |
| Rojas-Serrano et al. | Examples of adverse events are as follows: abdominal pain, anorexia, chest pain, constipation, diarrhea, dizziness, fatigue, gastroesophageal reflux, headache, nausea, paresthesia, rash, and throat tightness. Side effects include stomach, diarrhea, neurologic, headache, skin, palpitation, sleep disturbance, tinnitus, vision, allergic reaction, myalgia, bloody nose, appetite change, joint pain, low energy, mouth ulcers, yeast infection, dry mouth, and other. |
| McKinnon et al. (WHIP) | Covid-19 related symptoms, covid-19 clinical disease and medication adverse effects including gastrointestinal disorders, nervous system disorders, respiratory, thoracic and mediastinal disorders, general disorders and administration site conditions, cardiac disorders, musculoskeletal and connective tissue disorders, psychiatric disorders, skin and subcutaneous tissue disorders, ear and labyrinth disorders, and eye disorders. |
| Vijayaraghavan et al. | Adverse events listed in each category at the participant level were categorized as cardiac, gastro-intestinal, headache, and Qtc prolongation. |
| Polo et al. (EPICOS) | Adverse events were classified by organ system and included: gastrointestinal disorders, blood and lymphatic system disorders, cardiac disorders, ear and labyrinth disorders, eye disorder, general disorders, immune system disorder, infections, injuries, investigations, metabolism and nutrition disorders, musculoskeletal/connective tissue disorders, nervous system disorders, psychiatric disorders, renal and urinary disorders, reproductive system disorders, respiratory disorders, skin disorders and vascular disorders. |
| Llanos-Cuentas et al. | Adverse events from grade 1 to grade 3 and above. Note that the Llanos-Cuentas et al. study did report the number of adverse events (not participants) in the HCQ group only. Due to limited information, it was excluded from the meta-analysis with the adverse event outcome. |
| Grau-Pujol et al. | Adverse events included: general symptoms (fever, chills, sweating, malaise, myalgia, arthralgia), gastrointestinal symptoms (nausea, abdominal pain, diarrhea, dysgeusia), dermatological symptoms (itching, rash), respiratory symptoms (rhinorrhea, sore throat / odynophagia, cough, pleuritic pain, dyspnea), neurologic symptoms (headache, visual disturbances), and cardiovascular symptoms. Events were graded mild, moderate and severe. |
| Syed et al. | Syed et al. report the number of patients in each group who experienced adverse events, but did not report what the events were. Due to limited information, it was excluded from the meta-analysis with the adverse event outcome. |

eTable 5. Baseline characteristics with additional variables and detailed information. Sample mean and standard deviation (in parenthesis) are reported for continuous variables, and the number of participants and proportion (in parenthesis) are reported for binary or categorical variables.

| | | Naggie et al. (HERO-HCQ) | | Abella et al. (PATCH) | | Rajasingham et al. (MN-COVID-PREP) | | Rojas-Serrano et al. | | McKinnon et al. (WHIP) | |
|--------------------|---|-----------------------------|--------------------|--------------------------|-------------------------|---------------------------------------|--------------------------|-----------------------------|-------------------------------|--|-------------|
| | | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo |
| N (randomization) | | 683 | 676 | 66 | 66 | 989 | 494 | 62 | 65 | 387 | 191 |
| Age | | 44.2 (11.9) | 43.1 (11.2) | 31 (20-66) ³ | 34 (23-62) ³ | 41.5 (35, 49) ³ | 40 (34, 48) ³ | 31.0 (26.4-39) ⁴ | 31.9 (27.2-43.7) ⁴ | 45.7 (11.6); 44.9 (11.4) ² | 44.1 (12.7) |
| Female | | 442 (64.7%) | 446 (66.0%) | 54 (82%) | 37 (56%) | 519 (52.5%) | 241 (48.8%) | 29 (42.6%) | 42 (64.6%) | 220 (57%) | 114 (60%) |
| BMI (kg/m^2) | | 28.3 (6.3) | 28.6 (6.7) | 26 (19-37) ⁵ | 26 (20-50) ⁵ | | | 26.7 (3.9) | 27.2 (4.6) | | |
| Current smoker | | | | 0 (0%) | 0 (0%) | 38 (3.84%) | 13 (2.6%) | 20 (32.2%) ⁶ | 23 (35.4%) ⁶ | | |
| Race/ Ethnicity | White | 624 (91.4%) | 610 (90.2%) | 55 (83%) | 54 (82%) | 852 (86.1%) | 419 (84.8%) | | | 334 (86%) | 161 (84%) |
| | Asian | | | 7 (11%) | 7 (11%) | 46 (4.7%) | 29 (5.9%) | | | 23 (6%) | 15 (8%) |
| | African American | 18 (2.6%) | 23 (3.4%) | 3 (4%) | 1 (2%) | 10 (1.0%) | 10 (2.0%) | | | 15 (4%) | 9 (5%) |
| | Hispanic | 39 (5.7%) | 40 (5.9%) | 0 (0%) | 2 (3%) | 40 (4.0%) | 18 (3.6%) | | | 11 (3%) | 7 (4%) |
| Comorb idities | Asthma | 58 (8.5%) | 77 (11.4%) | 9 (14%) | 14 (21%) | 91 (9.2%) | 59 (11.9%) | | | | |
| | Diabetes | 20 (2.9%) | 35 (5.2%) | 1 (2%) | 3 (5%) | 36 (3.6%) | 14 (2.8%) | | | | |
| | Hypertension | 99 (14.5%) | 99 (14.6%) | 3 (5%) | 14 (21%) | 145 (14.7%) | 60 (12.1%) | | | | |
| | None | | | 54 (82%) | 40 (61%) | 646 (65.3%) | 336 (68.0%) | 53 (85.5%) | 58 (89.2%) | | |
| Practice Location | Emergency Department | 96 (14.1%) | 94 (13.9%) | 38 (58%) | 36 (55%) | 417 (42.2%) | 190 (38.5%) | | | 48 (12%) | 19 (10%) |
| | Internal Medicine ward | | | 17 (26%) | 18 (27%) | 98 (9.9%) | 56 (11.3%) | | | 31 (8%) | 20 (10%) |
| | ICU/anesthesia | | | 6 (9%) | 6 (9%) | | | | | | |
| | Labor and delivery | | | 5 (7%) | 6 (9%) | | | | | | |
| | Ambulance | 66 (9.7%) | 63 (9.3%) | | | 73 (7.4%) | 45 (9.1%) | | | | |
| | Congregate care setting | | | | | 46 (4.7%) | 20 (4.0%) | | | | |
| | ICU | 48 (7.0%) | 59 (8.7%) | | | 184 (18.6%) | 85 (17.2%) | | | 37 (10%) | 23 (12%) |
| | Operating room | | | | | 103 (10.4%) | 75 (15.2%) | | | | |
| | EMS, Fire and Police | | | | | | | | | 32 (8%) | 16 (8%) |
| | First Responders | | | | | | | | | | |
| Occupation | Nurse | 186/677 (27.5%) | 167/668 (25.0%) | 46 (70%) | 42 (64%) | | | | | | |
| | Physician | 143/677 (21.1%) | 144/668 (21.6%) | 11 (17%) | 16 (24%) | | | | | | |
| | Certified Nurse Assistant | | | 2 (3%) | 2 (3%) | | | | | | |
| | ED Technician | | | 3 (4%) | 1 (2%) | | | | | | |
| | Respiratory therapist | 15/677 (2.2%) | 18/668 (2.7%) | 3 (4%) | 5 (7%) | | | | | | |
| | Nurse or Physician | | | | | | | 31 (50%) | 33 (50.8%) | | |
| | Emergency Medicine Provider | | | | | 407 (41.1%) | 190 (38.5%) | | | | |
| | ICU provider | | | | | 160 (16.2%) | 83 (16.8%) | | | | |
| | Anesthesia/ENT | | | | | 178 (18.0%) | 105 (21.3%) | | | | |
| | HCW in COVID unit | | | | | 76 (7.7%) | 29 (5.9%) | | | | |
| | Healthcare worker in congregated care setting | | | | | 11 (1.1%) | 4 (0.8%) | | | | |
| | First responder | | | | | 115 (11.6%) | 65 (13.2%) | | | | |

| | | Vijayaraghavan et al. | | Polo et al. (EPICOS) | | Llanos-Cuentas et al. | | Grau-Pujol et al. | | Syed et al. | |
|--------------------------|--|-----------------------|-------------|-------------------------|-------------|-----------------------|--------------|-------------------|-------------|------------------|-------------|
| | | HCQ | Placebo | HCQ ² | Placebo | HCQ | Placebo | HCQ | Placebo | HCQ ¹ | Placebo |
| | N (randomization) | 213 | 203 | 231 | 223 | 36 | 32 | 142 | 127 | 154 | 46 |
| | Age | 32.3 (9.65) | 31.8 (8.63) | 38 (18-65) | 38 (18,65) | 39.14 (1.53) | 39.28 (1.72) | 39.6 (11.2) | 40.3 (12.8) | 30.25 (NA) | 31.9 (9.13) |
| Female | | 100 (46.9%) | 97 (47.8%) | 149 (64.5%) | 143 (64.1%) | 20 (55.6%) | 20 (62.5%) | 104 (73.2%) | 93 (73.2%) | 68 (44.1%) | 23 (50%) |
| BMI (kg/m ²) | | | | | | | | | | | |
| Current smoker | | 8 (3.8%) | 9 (4.4%) | | | | | 21 (14.9%) | 17 (13.8%) | 19 (12.3%) | 7 (15.2%) |
| Race/ Ethnicity | White | | | | | | | | | | |
| | Asian | | | | | | | | | | |
| | African American | | | | | | | | | | |
| | Hispanic | | | | | | | | | | |
| Comorb idities | Asthma | 0 (0%) | 0 (0%) | 20 (8.7%) | 9 (4.0%) | 3 (8.3%) | 4 (12.5%) | 5 (3.5%) | 2 (1.6%) | | |
| | Diabetes | 7 (3.3%) | 3 (1.5%) | 1 (0.4%) | 3 (1.3%) | 1 (2.8%) | 0 (0%) | 0 (0%) | 1 (0.8%) | 4 (2.6%) | 3 (6.5%) |
| | Hypertension | 2 (0.9%) | 3 (1.5%) | 4 (1.7%) | 19 (8.5%) | 3 (8.3%) | 2 (6.3%) | 2 (1.4%) | 3 (2.4%) | 7 (4.5%) | 2 (4.3%) |
| | None | | | | | | | | | | |
| Practice Location | Emergency Department | 26 (12.2%) | 18 (8.9%) | 20 (8.7%) | 21 (9.4%) | | | | | | |
| | Internal Medicine ward | 130 (64%) | 130 (61%) | | | | | | | | |
| | ICU/anesthesia | | | | | | | | | | |
| | Labor and delivery | | | | | | | | | | |
| | Ambulance | | | 0 (0%) | 0 (0%) | | | | | | |
| | Congregate care setting | | | | | | | | | | |
| | ICU | 53 (24.9%) | 53 (26.1%) | 17 (7.4%) | 13 (5.8%) | | | | | | |
| | Operating room | | | | | | | | | | |
| | EMS, Fire and Police First Responders | | | | | | | | | | |
| Occupation | Nurse | 67 (31.5%) | 68 (33.5%) | 67 (29.0%) | 72 (32.3%) | 6 (16.7%) | 5 (15.6%) | 35 (27.8%) | 40 (28.2%) | 20 (13.0%) | 9 (19.6%) |
| | Physician | 34 (16%) | 31 (15.3%) | 74 (32%) | 66 (29.6%) | 23 (63.9%) | 16 (50%) | 67 (47.2%) | 53 (42.1%) | 118 (76.6%) | 25 (54.3%) |
| | Certified Nurse Assistant | | | | | 1 (2.8%) | 0 (0%) | 12 (8.5%) | 12 (9.5%) | | |
| | ED Technician | | | | | | | | | | |
| | Respiratory therapist | | | | | | | | | | |
| | Nurse or Physician | | | | | | | | | | |
| | Emergency Medicine Provider | | | | | | | | | 2 (1.3%) | 0 (0%) |
| | ICU provider | | | | | | | | | | |
| | Anesthesia/ENT | | | | | | | | | | |
| | HCW in COVID unit | | | | | | | | | | |
| | Healthcare worker in congregate care setting | | | | | | | | | | |
| | First responder | | | | | | | | | 2 (1.3%) | 0 (0%) |

HCQ=Hydroxychloroquine ; ITT= Intention-to-treat ; BMI=Body mass index ; ICU=Intensive care unit; ED=Emergency department ; ENT=Ear, nose, throat ; HCW=Healthcare worker

¹ More than one HCQ groups with different doses are lumped.

² The Polo et al. study randomized participants to four treatment groups, and the HCQ and control groups are used in our meta-analysis.

³ Median (range)

⁴ Median (IQR)

⁵ Mean (range)

⁶ Current or previous smoker

eTable 6. Results of outcome measures in trials included in the meta-analysis. Sample size and the number of participants who had each outcome are reported with proportions (%) in parentheses.

| | Treatment | N (ITT) | Confirmed COVID-19 | Suspected with COVID compatible symptoms | Adverse event ² |
|------------------------------------|------------------|---------|--------------------|--|----------------------------|
| Naggie et al. (HERO-HCQ) | HCQ | 683 | 3 (0.4) | 38 (5.6) | 16 (2.3) |
| | Placebo | 676 | 6 (0.9) | 47 (7.0) | 13 (1.9) |
| Abella et al. (PATCH) | HCQ | 64 | 4 (6.3) | | 29 (45.3) |
| | Placebo | 61 | 4 (6.6) | | 17 (27.9) |
| Rajasingham et al. (MN-COVID-PREP) | HCQ ¹ | 989 | 11 (1.1) | 47 (4.8) | 316 (32.0) |
| | Placebo | 494 | 6 (1.2) | 33 (6.7) | 100 (20.2) |
| Rojas-Serrano et al. | HCQ | 62 | 1 (1.6) | | 32 (51.6) |
| | Placebo | 65 | 6 (9.2) | | 38 (58.5) |
| McKinnon et al. (WHIP) | HCQ ¹ | 387 | 2 (0.5) | 22 (5.7) | 192 (49.6) |
| | Placebo | 191 | 2 (1.0) | 13 (6.8) | 85 (44.5) |
| Vijayaraghavan et al. | HCQ | 211 | 11 (5.2) | 12 (5.7) | 21 (10.0) |
| | Placebo | 203 | 12 (5.9) | 12 (5.9) | 14 (6.9) |
| Polo et al. (EPICOS) | HCQ | 224 | 21 (9.4) | | 100 (44.6) |
| | Placebo | 211 | 23 (10.9) | | 94 (44.5) |
| Llanos-Cuentas et al. | HCQ | 34 | 5 (14.7) | | |
| | Placebo | 31 | 3 (9.7) | | |
| Grau-Pujol et al. | HCQ | 137 | 1 (0.7) | 3 (2.2) | 53 (38.7) |
| | Placebo | 116 | 1 (0.9) | 3 (2.6) | 42 (36.2) |
| Syed et al. | HCQ ¹ | 154 | 42 (27.3) | | 9 (5.8) |
| | Placebo | 46 | 7 (15.2) | | 1 (2.2) |


HCQ= Hydroxychloroquine; ITT= Intention-to-treat; AE=Adverse event ; COVID-RS=COVID-19 related symptoms ; Vit C= Vitamin C

¹ More than one HCQ groups with different doses are lumped.


² Number of patients with any adverse events

eFigure. Funnel plots for the three outcomes


(a) Lab-confirmed positive COVID-19



(b) Suspected COVID-19



(c) Adverse events



eTable 7. GRADE summary of findings table

| Outcomes | No of participants (studies) Follow up | Quality of the evidence (GRADE) | Odds ratio (95% Confidence Interval) |
|------------------------------------|--|--|---|
| Lab-confirmed positive COVID-19 | 5039 (10 studies) From 28 days to 180 days | ⊕⊕⊕⊖ Moderate ¹ due to imprecision | 0.92 (0.58, 1.37) |
| Suspected COVID-19 | 4087 (5 studies) From 56 days to 180 days | ⊕⊕⊕⊖ Moderate ¹ due to imprecision | 0.78 (0.57, 1.10) |
| Adverse events | 4979 (9 studies) From 56 days to 180 days | ⊕⊕⊕⊖ Moderate ² due to imprecision | 1.35 (1.03, 1.73) |

¹95% confidence interval includes effect suggesting benefit as well as no benefit.

²Although the 95% confidence interval includes an effect suggesting no benefit, we decided to downgrade it by one level because the lower limit is close to the null.

GRADE Working Group grades of evidence is available here:

<https://gdt.gradeapro.org/app/handbook/handbook.html>



PRISMA 2020 Checklist

| Section and Topic | Item # | Checklist item | Location where item is reported |
|-------------------------------|--------|--|---------------------------------|
| TITLE | | | |
| Title | 1 | Identify the report as a systematic review. | 1 |
| ABSTRACT | | | |
| Abstract | 2 | See the PRISMA 2020 for Abstracts checklist. | 4 |
| INTRODUCTION | | | |
| Rationale | 3 | Describe the rationale for the review in the context of existing knowledge. | 5-6 |
| Objectives | 4 | Provide an explicit statement of the objective(s) or question(s) the review addresses. | 6 |
| METHODS | | | |
| Eligibility criteria | 5 | Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses. | 7 |
| Information sources | 6 | Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted. | 7 |
| Search strategy | 7 | Present the full search strategies for all databases, registers and websites, including any filters and limits used. | 7 |
| Selection process | 8 | Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process. | 7 |
| Data collection process | 9 | Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process. | 7 |
| Data items | 10a | List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect. | 8 |
| | 10b | List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information. | 8 |
| Study risk of bias assessment | 11 | Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process. | 8 |
| Effect measures | 12 | Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results. | 9 |
| Synthesis methods | 13a | Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)). | Supplement |
| | 13b | Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions. | 9 |
| | 13c | Describe any methods used to tabulate or visually display results of individual studies and syntheses. | 8-9 |
| | 13d | Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used. | 10 |
| | 13e | Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression). | 10 |
| | 13f | Describe any sensitivity analyses conducted to assess robustness of the synthesized results. | 10 |
| Reporting bias assessment | 14 | Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases). | 9 |
| Certainty assessment | 15 | Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome. | 9 |



PRISMA 2020 Checklist

| Section and Topic | Item # | Checklist item | Location where item is reported |
|--|--------|--|---------------------------------|
| RESULTS | | | |
| Study selection | 16a | Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram. | 11 |
| | 16b | Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded. | 11-12 |
| Study characteristics | 17 | Cite each included study and present its characteristics. | 8-9, Supplement |
| Risk of bias in studies | 18 | Present assessments of risk of bias for each included study. | Supplement |
| Results of individual studies | 19 | For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots. | Supplement |
| Results of syntheses | 20a | For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies. | Supplement |
| | 20b | Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect. | 11-13 |
| | 20c | Present results of all investigations of possible causes of heterogeneity among study results. | 11-13 |
| | 20d | Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results. | 11-13 |
| Reporting biases | 21 | Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed. | Supplement |
| Certainty of evidence | 22 | Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed. | Supplement |
| DISCUSSION | | | |
| Discussion | 23a | Provide a general interpretation of the results in the context of other evidence. | 14 |
| | 23b | Discuss any limitations of the evidence included in the review. | 16 |
| | 23c | Discuss any limitations of the review processes used. | 16 |
| | 23d | Discuss implications of the results for practice, policy, and future research. | 16 |
| OTHER INFORMATION | | | |
| Registration and protocol | 24a | Provide registration information for the review, including register name and registration number, or state that the review was not registered. | Supplement |
| | 24b | Indicate where the review protocol can be accessed, or state that a protocol was not prepared. | 7 |
| | 24c | Describe and explain any amendments to information provided at registration or in the protocol. | 7 |
| Support | 25 | Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review. | 17 |
| Competing interests | 26 | Declare any competing interests of review authors. | 17 |
| Availability of data, code and other materials | 27 | Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review. | Supplement |